

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Banaji, Murad ORCID: <https://orcid.org/0000-0002-4983-0377> and Rutherford, Carrie (2010)
P-matrices and signed digraphs. *Discrete Mathematics*, 311 (4) . pp. 295-301. ISSN
0012-365X [Article] (doi:10.1016/j.disc.2010.10.018)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/20584/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

P -matrices and signed digraphs

Murad Banaji^{a,*}, Carrie Rutherford^b

^a*Department of Mathematics, University of Portsmouth, Lion Gate Building, Lion Terrace, Portsmouth, Hampshire PO1 3HF, UK.*

^b*Department of Business Studies, London South Bank University, 103 Borough Road, London SE1 0AA, UK*

Abstract

We associate a signed digraph with a list of matrices whose dimensions permit them to be multiplied, and whose product is square. Cycles in this graph have a parity, that is, they are either even (termed e-cycles) or odd (termed o-cycles). The absence of e-cycles in the graph is shown to imply that the matrix product is a P_0 -matrix, i.e., all of its principal minors are nonnegative. Conversely, the presence of an e-cycle is shown to imply that there exists a list of matrices associated with the graph whose product fails to be a P_0 -matrix. The results generalise a number of previous results relating P - and P_0 -matrices to graphs.

Keywords: P -matrix, matrix factorisation, signed digraph, SR graph

1. Introduction and statement of the main results

P -matrices are square matrices, all of whose principal minors are positive. P_0 -matrices [1] are square matrices all of whose principal minors are nonnegative, i.e., matrices in the closure of the P -matrices. We will be interested in real P_0 -matrices. Amongst other areas, P - and P_0 -matrices play important roles in the study of linear complementarity problems [2, 3] and in population biology [4]. A well-known result of Gale and Nikaido [5], whose extensions and corollaries are discussed in [6], states that a differentiable function on a rectangular domain in \mathbb{R}^n with P -matrix Jacobian is injective. This result has a number of practical applications – see for example [7, 8].

The results of Gale and Nikaido have various graph-theoretic implications, explored in [9, 10, 11, 12] for example. The key results in these papers involve determining sufficient graph-theoretic conditions for a set of matrices to be P_0 -matrices, followed by additional “nondegeneracy” conditions which guarantee that they are in fact P -matrices. A feature of this work is that matrices which arise in applications often have a natural *product* structure. For example, the following question is shown to be pertinent to problems in biology and chemistry in [10, 11]: given a set \mathcal{M} of $n \times m$ matrices, and a set \mathcal{N} of $m \times n$ matrices, is

*Corresponding author: murad.banaji@port.ac.uk.

$\{MN : M \in \mathcal{M}, N \in \mathcal{N}\}$ a subset of the P_0 -matrices? Here, a general result is developed, of which some of these previous results become corollaries.

Let k be an arbitrary positive integer. Throughout this paper, a subscript or superscript j assumed to belong to $\{0, \dots, k-1\}$ should be read as $j \bmod k$. Let n_0, \dots, n_{k-1} be positive integers. For each $j \in \{0, \dots, k-1\}$, let $A^{(j)}$ be an $n_j \times n_{j+1}$ matrix, and define the $n_0 \times n_0$ matrix $A = A^{(0)}A^{(1)} \dots A^{(k-1)}$. We will associate with $[A^{(0)}, A^{(1)}, \dots, A^{(k-1)}]$ a signed digraph $G_{A^{(0)} \dots A^{(k-1)}}$, which will belong to a category of graphs termed signed $(k, \{1\})$ -BC digraphs, to be defined below. The structure of these graphs will imply that all its cycles have length which is a multiple of k .

Given a cycle C with kr_1 edges, of which r_2 have negative sign, we define C to be an **e-cycle** if $(-1)^{r_1+r_2} = 1$ and an **o-cycle** otherwise. A signed $(k, \{1\})$ -BC digraph containing no e-cycles will be termed “e-cycle-free”. The first main theorem in this paper is:

Theorem 1. *If $G = G_{A^{(0)} \dots A^{(k-1)}}$ is e-cycle-free then $A = A^{(0)}A^{(1)} \dots A^{(k-1)}$ is a P_0 -matrix.*

A matrix M determines the qualitative class $\mathcal{Q}(M)$ [13] consisting of all matrices with the same sign pattern as M . Explicitly, $\mathcal{Q}(M)$ consists of all matrices X with the same dimensions as M , and satisfying $M_{ij} > 0 \Rightarrow X_{ij} > 0$, $M_{ij} < 0 \Rightarrow X_{ij} < 0$ and $M_{ij} = 0 \Rightarrow X_{ij} = 0$. Given two matrices M and N of dimensions such that they can be multiplied, we write

$$\mathcal{Q}(M)\mathcal{Q}(N) = \{M'N' \mid M' \in \mathcal{Q}(M), N' \in \mathcal{Q}(N)\}.$$

This definition extends naturally to any ordered set of multiplicable matrices. Note that $\mathcal{Q}(M)\mathcal{Q}(N)$ is in general different from $\mathcal{Q}(MN)$, and is often not a subset of any qualitative class. The second main theorem in this paper is:

Theorem 2. *All matrices in $\mathcal{Q}(A^{(0)})\mathcal{Q}(A^{(1)}) \dots \mathcal{Q}(A^{(k-1)})$ are P_0 -matrices if and only if $G_{A^{(0)}A^{(1)} \dots A^{(k-1)}}$ is e-cycle-free.*

2. Signed $(k, \{1\})$ -BC digraphs

Consider a digraph G with vertex set $V(G)$ and edge set $E(G)$. Let S be any set of integers. G will be termed “ (k, S) -block circulant”, abbreviated to (k, S) -BC, if

1. $V(G)$ is partitioned into k sets $\{V_0, V_1, \dots, V_{k-1}\}$.
2. For $k \geq 2$, if $(j-i) \bmod k \notin S$, then there is no (directed) edge from a vertex in V_i to a vertex in V_j .

Every digraph is trivially a $(1, S)$ -BC digraph for arbitrary S . When $k \geq 2$, (k, S) -BC digraphs are a generalisation of circulant digraphs ([14, 15] for example). Note, however, that vertices in a (k, S) -BC digraph may have arbitrary outdegree and indegree. Here, only the special case $S = \{1\}$ concerns us. In a $(k, \{1\})$ -BC digraph, a (directed) path from a vertex in V_i to a vertex in V_j must

include a vertex from each $V_{j'}$, $j' \neq j$. It follows that all cycles in a $(k, \{1\})$ -BC digraph have length which is a multiple of k .

Remark. Although, for $k \geq 2$, $(k, \{1\})$ -BC digraphs are k -colourable, k is not in general the chromatic number of G : for example, a $(2r, \{1\})$ -BC digraph with nonempty edge-set is in fact bipartite.

Let V_j contain n_j vertices. Assume some ordering on these vertices and let V_j^i ($i \in \{1, \dots, n_j\}$) refer to the i th vertex in V_j . As usual, an edge (v, \tilde{v}) refers to the edge directed from v to \tilde{v} .

A digraph G is signed if there is a function $\text{sign} : E(G) \rightarrow \{-1, +1\}$. A signed $(k, \{1\})$ -BC digraph $G = G_{A^{(0)} \dots A^{(k-1)}}$ is associated with a list of matrices $[A^{(0)}, \dots, A^{(k-1)}]$ as above in a very simple way: (i) for each $j = 0, \dots, k-1$, $|V_j| = n_j$, (ii) there exists an edge (V_j^r, V_{j+1}^s) in G if and only if $(A^{(j)})_{rs} \neq 0$, and (iii) the edge (V_j^r, V_{j+1}^s) takes the sign of $(A^{(j)})_{rs}$. Note that entries in the matrices $A^{(j)}$ are in one-to-one correspondence with edges in G , and that the sign-pattern of $A^{(j)}$ is in fact a block in the (signed) adjacency matrix of G .

Example. As an example consider the matrices:

$$A^{(0)} = \begin{pmatrix} a & -b & -c \\ d & 0 & e \end{pmatrix}, \quad A^{(1)} = \begin{pmatrix} f & 0 \\ -g & 0 \\ 0 & h \end{pmatrix}, \quad A^{(2)} = \begin{pmatrix} w & x \\ -y & z \end{pmatrix}, \quad (1)$$

where $a, b, c, d, e, f, g, h, w, x, y$ and z are arbitrary positive real numbers. Associated with the product $A^{(0)}A^{(1)}A^{(2)}$ is the signed $(3, \{1\})$ -BC digraph shown in Figure 1.

Defining $\mathbf{0}_{m \times n}$ to be the $m \times n$ zero matrix, the signed adjacency matrix of the graph G in Figure 1 has block structure:

$$\begin{pmatrix} \mathbf{0}_{2 \times 2} & \begin{pmatrix} 1 & -1 & -1 \\ 1 & 0 & 1 \end{pmatrix} & \mathbf{0}_{2 \times 2} \\ \mathbf{0}_{3 \times 2} & \mathbf{0}_{3 \times 3} & \begin{pmatrix} 1 & 0 \\ -1 & 0 \\ 0 & 1 \end{pmatrix} \\ \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} & \mathbf{0}_{2 \times 3} & \mathbf{0}_{2 \times 2} \end{pmatrix}.$$

It can be seen immediately that each block is simply the sign-pattern of $A^{(0)}$, $A^{(1)}$ or $A^{(2)}$.

Although the graph in Figure 1 has a number of cycles (both of length 3 and of length 6), all of these can be computed to be o-cycles, and so, by Theorem 1, the product $A^{(0)}A^{(1)}A^{(2)}$ is a P_0 -matrix. This is true whatever the magnitudes of the entries in the matrices. Clearly, given isomorphic signed digraphs $G_1 \cong G_2$, G_1 is e-cycle-free if and only if G_2 is e-cycle-free. Since

$$G_{A^{(1)}A^{(2)}A^{(0)}} \cong G_{A^{(2)}A^{(0)}A^{(1)}} \cong G_{A^{(0)}A^{(1)}A^{(2)}},$$

$A^{(1)}A^{(2)}A^{(0)}$ and $A^{(2)}A^{(0)}A^{(1)}$ are also P_0 -matrices.




Figure 1: The signed $(3, \{1\})$ -BC digraph corresponding to the product of the three matrices $A^{(0)}A^{(1)}A^{(2)}$ in Eq. 1. The graph has been laid out to emphasise its circulant structure. The vertices in V_0, V_1 and V_2 are represented as filled circles, open circles and boxes respectively. Negative edges are represented as dashed lines while positive edges are bold lines. Labels on each edge represent the absolute values of the corresponding entries in the matrices, and are not strictly part of the graph: they have been added to indicate the correspondence between edges and matrix entries.

3. Preliminaries needed for the proofs

Permutations. Consider an ordered set $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_r]$, and a bijection $\beta : \alpha \rightarrow \alpha$. Defining $\beta_i \equiv \beta(\alpha_i)$, the ordered set $[\beta_1, \beta_2, \dots, \beta_r]$ will also be referred to as β . It will always be clear from context whether an object referred to is a bijection or an ordered set.

Any permutation β has a sign $\sigma(\beta)$, i.e., $\sigma(\beta) = +1$ if β is an even permutation and $\sigma(\beta) = -1$ otherwise. Given two permutations β and β' , $\sigma(\beta\beta') = \sigma(\beta)\sigma(\beta')$, implying that $\sigma(\beta) = \sigma(\beta^{-1})$. Note the following elementary result about the sign of a permutation.

Lemma 3. Consider a permutation β of a finite set of size r . Write β as a product of disjoint cycles, C_1, \dots, C_s ($1 \leq s \leq r$), including trivial cycles. Then

$$\sigma(\beta) = (-1)^{r-s}$$

i.e., β is even (resp. odd) if the total number of elements in β minus the total number of cycles in its decomposition is even (resp. odd).

PROOF. This follows immediately by noting that the sign of a product of cycles is the product of the cycles' signs, and that a k -cycle is an even permutation if k is odd and vice versa.

From here on $\alpha^{(j)}$ will always refer to a nonempty subset of $\{1, \dots, n_j\}$, and will be assumed to have the natural ordering. $\alpha_m^{(j)}$ will refer to the m th element in $\alpha^{(j)}$ so that $\alpha_1^{(j)} < \alpha_2^{(j)} < \alpha_3^{(j)} < \dots$. Given some $\alpha^{(j)}$, define $V_j^{\alpha^{(j)}} = \{V_j^k \mid k \in \alpha^{(j)}\} \subseteq V_j$. $\beta^{(j)}$ will refer to a permutation of $\alpha^{(j)}$. Given the one-to-one correspondence between the elements in $\alpha^{(j)}$, and vertices in $V_j^{\alpha^{(j)}}$, $\beta^{(j)}$ can equally be regarded as a permutation on $V_j^{\alpha^{(j)}}$.

Now consider some sequence $(\alpha^{(0)}, \dots, \alpha^{(k-1)})$, such that $|\alpha^{(i)}| = |\alpha^{(j)}|$ for each i, j , and a corresponding sequence of permutations $(\beta^{(0)}, \dots, \beta^{(k-1)})$. Define $R = [1, 2, \dots, |\alpha^{(0)}|]$, and define the bijections $\iota_j : V_j^{\alpha^{(j)}} \rightarrow R$ by $\iota_j(V_j^{\alpha_m^{(j)}}) = m$. In other words, ι_j associates with each vertex in $V_j^{\alpha^{(j)}}$ its order. $\beta^{(j)}$ then induces the bijection $\tilde{\beta}^{(j)} : R \rightarrow R$ defined by $\tilde{\beta}^{(j)} = \iota_j \circ \beta^{(j)} \circ \iota_j^{-1}$. Further, define the bijections $\phi_j : V_j^{\alpha^{(j)}} \rightarrow V_{j+1}^{\alpha^{(j+1)}}$ by $\phi_j = \iota_{j+1}^{-1} \circ \tilde{\beta}^{(j+1)} \circ \iota_j$, i.e. $\phi_j(V_j^{\alpha_m^{(j)}}) = V_{j+1}^{\beta_m^{(j+1)}}$. Equivalently, $\phi_j = \beta^{(j+1)} \circ \iota_{j+1}^{-1} \circ \iota_j$. Figure 2 illustrates all of these relationships. Note that in the special case $k = 1$, $\phi_j = \beta^{(j)}$.




Figure 2: The commutative diagram which encapsulates the relationships $\tilde{\beta}^{(j)} = \iota_j \circ \beta^{(j)} \circ \iota_j^{-1}$, $\phi_j = \beta^{(j+1)} \circ \iota_{j+1}^{-1} \circ \iota_j$, and $\phi_j = \iota_{j+1}^{-1} \circ \tilde{\beta}^{(j+1)} \circ \iota_j$.

4. Proof of Theorem 1 and its immediate consequences

The following notation is used. Given an $r \times s$ matrix M , and two (nonempty) ordered sets $\gamma \subseteq \{1, \dots, r\}$ and $\delta \subseteq \{1, \dots, s\}$, then $M(\gamma|\delta)$ is the submatrix of M with rows indexed by γ and columns indexed by δ . If $|\gamma| = |\delta|$, then $M[\gamma|\delta] \equiv \det(M(\gamma|\delta))$. We write $M[\gamma]$ as shorthand for $M[\gamma|\gamma]$. If γ and δ are of equal size, then $M_{\gamma,\delta}$ will refer to $\prod_{i=1}^{|\gamma|} M_{\gamma_i,\delta_i}$.

PROOF OF THEOREM 1. The cases $k = 1$ and $k \geq 2$ are conceptually similar:

however in order to avoid notational difficulties, they are presented separately.

Case 1: $k = 1$. Let $V = V(G)$. Choose and fix some nonempty $\alpha \subseteq \{1, \dots, n\}$, and consider the minor $A[\alpha]$. Given a permutation β of α , define T , a term in $A[\alpha]$, by:

$$T = \sigma(\beta)A_{\alpha,\beta}.$$

Suppose that $T \neq 0$. Since nonzero entries in A are in one-to-one correspondence with edges in G , corresponding to T is an edge-set $E \subseteq E(G)$ defined as follows: the edge (V^r, V^s) is in E if and only if $(r, s) = (\alpha_m, \beta_m)$ for some integer m . Equivalently, the edge (V^r, V^s) is in E if and only if $V^s = \beta(V^r)$.

The set of endpoints of edges in E is precisely V^α . Consider the vertex V^{α_m} . Then the incoming edge $(\beta^{-1}(V^{\alpha_m}), V^{\alpha_m})$ and the outgoing edge $(V^{\alpha_m}, \beta(V^{\alpha_m}))$ are the only edges from E incident on V^{α_m} . If $\beta(V^{\alpha_m}) = V^{\alpha_m}$, then these edges coincide and in fact there is a loop at V^{α_m} . Otherwise, the edges are distinct. In either case, E , regarded as a subgraph of G , consists of vertex- and edge-disjoint cycles.

Suppose $\beta^m(v) = v$ for some vertex v , but $\beta^q(v) \neq v$ for $q < m$. Then the vertex v lies on a cycle in E of length m . Thus, decomposing β as a product of disjoint cycles (including trivial cycles), these cycles are in one-to-one correspondence with cycles – in the graph-theoretic sense – in E . Trivial cycles correspond to loops. Assume that there are N such cycles. By Lemma 3, $\sigma(\beta) = (-1)^{|\alpha| - N}$.

Returning to the term T in the expansion of $A[\alpha]$,

$$\text{sign}(T) = \sigma(\beta)\text{sign}(A_{\alpha,\beta}) \tag{2}$$

Consider a cycle C in E including r_1 edges of which r_2 are negative, and define $\text{sign}(C) = (-1)^{r_2}$ (i.e. $\text{sign}(c)$ is the product of signs of edges in C). If C is an e-cycle, then $(-1)^{r_1+r_2} = 1$, and so $\text{sign}(C) = (-1)^{r_1}$. Similarly if C is an o-cycle, then $\text{sign}(C) = (-1)^{r_1+1}$. Decompose E into disjoint cycles, which comprise N_e e-cycles, and N_o o-cycles (so that $N = N_o + N_e$). Let θ be the total number of edges in o-cycles and θ_e the total number of edges in e-cycles, so that $\theta_o + \theta_e = |\alpha|$ (since there are $|\alpha|$ edges in E). Taking the product of signs of edges in E over e-cycles and o-cycles separately gives

$$\text{sign}(A_{\alpha,\beta}) = (-1)^{\theta_o}(-1)^{\theta_e+N_o} = (-1)^{|\alpha|+N_o}. \tag{3}$$

Since $\sigma(\beta) = (-1)^{|\alpha| - N}$, and $\text{sign}(A_{\alpha,\beta}) = (-1)^{|\alpha|+N_o}$, Eq. 2 gives:

$$\text{sign}(T) = (-1)^{|\alpha| - N}(-1)^{|\alpha|+N_o} = (-1)^{N_e}. \tag{4}$$

If G is e-cycle-free, then $N_e = 0$ in this expression, in which case $\text{sign}(T) = 1$. Since T is an arbitrary nonzero term in $A[\alpha]$, $A[\alpha] \geq 0$. Since α is an arbitrary nonempty subset of $\{1, \dots, n\}$, A is a P_0 -matrix.

Case 2: $k \geq 2$. Choose and fix some nonempty $\alpha^{(0)} \subseteq \{1, \dots, n_0\}$, and consider the minor $A[\alpha^{(0)}]$. Enumerate all lists $(\alpha^{(1)}, \dots, \alpha^{(k-1)})$ where for each $i = 1, \dots, k-1$, $\alpha^{(i)}$ is a nonempty subset of $\{1, \dots, n_i\}$ such that $|\alpha^{(i)}| = |\alpha^{(0)}|$. There may of course be no such subset for some i , and hence no such lists.

By the Cauchy-Binet formula [16] applied recursively,

$$A[\alpha^{(0)}] = \sum_{\substack{(\alpha^{(1)}, \dots, \alpha^{(k-1)}) \\ |\alpha^{(i)}| = |\alpha^{(0)}|}} \left(\prod_{j=0}^{k-1} A^{(j)}[\alpha^{(j)} | \alpha^{(j+1)}] \right).$$

The sum is over all possible lists $(\alpha^{(1)}, \dots, \alpha^{(k-1)})$ with $|\alpha^{(i)}| = |\alpha^{(0)}|$. Now choose and fix some particular choice $\alpha^{(1)}, \dots, \alpha^{(k-1)}$, and choose permutations $\beta^{(0)}, \dots, \beta^{(k-1)}$. For each j define T_j , a term in $A^{(j)}[\alpha^{(j)} | \alpha^{(j+1)}]$, by:

$$T_j = \sigma(\beta^{(j+1)}) A_{\alpha^{(j)}, \beta^{(j+1)}}^{(j)}.$$

Suppose that for each j , T_j is nonzero so that $T = \prod_j T_j \neq 0$. Note that T is then a nonzero term in the expansion of $A[\alpha^{(0)}]$. Since nonzero entries in the matrices $A^{(j)}$ are in one-to-one correspondence with edges in G , corresponding to T is an edge-set $E \subseteq E(G)$ defined as follows: the edge (V_j^r, V_{j+1}^s) is in E if and only if $(r, s) = (\alpha_m^{(j)}, \beta_m^{(j+1)})$ for some integer m . Equivalently, the edge (V_j^r, V_{j+1}^s) is in E if and only if $\phi_j(V_j^r) = V_{j+1}^s$.

The set of endpoints of edges in E is precisely $\bigcup_j V_j^{\alpha^{(j)}}$, and in fact each such vertex has exactly two edges from E incident on it, one incoming and one outgoing. For example, consider the vertex V_j^r , where $r \in \alpha^{(j)}$. Then the incoming edge $(\phi_{j-1}^{-1}(V_j^r), V_j^r)$ and the outgoing edge $(V_j^r, \phi_j(V_j^r))$ are distinct edges in E , and are, by the definition of E , the only two edges in E incident on V_j^r . As a consequence, E , regarded as a subgraph of G , consists of vertex- and edge-disjoint cycles.

Next, consider the bijection $\phi : V_0^{\alpha^{(0)}} \rightarrow V_0^{\alpha^{(0)}}$ defined by $\phi = \phi_{k-1} \circ \phi_{k-2} \circ \dots \circ \phi_0$. Suppose $\phi^m(v) = v$ for some vertex v , but $\phi^q(v) \neq v$ for $q < m$. Then the vertex v lies on a cycle in E of length km . Decomposing ϕ as a product of disjoint cycles (including trivial cycles), these cycles are in one-to-one correspondence with cycles – in the graph-theoretic sense – in E . Assume that there are N such cycles. By Lemma 3, $\sigma(\phi) = (-1)^{|\alpha^{(0)}| - N}$. Applying the relations $\phi_j = \iota_{j+1}^{-1} \circ \tilde{\beta}^{(j+1)} \circ \iota_j$ gives

$$\phi = \iota_0^{-1} \circ \tilde{\beta}^{(0)} \circ \tilde{\beta}^{(k-1)} \circ \dots \circ \tilde{\beta}^{(2)} \circ \tilde{\beta}^{(1)} \circ \iota_0$$

so that $\sigma(\phi) = \prod_{j=0}^{k-1} \sigma(\tilde{\beta}^{(j)}) = \prod_{j=0}^{k-1} \sigma(\beta^{(j)})$.

Returning to the term T in the expansion of $A[\alpha^{(0)}]$,

$$\text{sign}(T) = \prod_{j=0}^{k-1} \text{sign}(T_j) = \left(\prod_{j=0}^{k-1} \sigma(\beta^{(j+1)}) \right) \left(\prod_{j=0}^{k-1} \text{sign} \left(A_{\alpha^{(j)}, \beta^{(j+1)}}^{(j)} \right) \right). \quad (5)$$

The first term in this expression has already been determined: from above,

$$\prod_{j=0}^{k-1} \sigma(\beta^{(j+1)}) = \prod_{j=0}^{k-1} \sigma(\beta^{(j)}) = \sigma(\phi) = (-1)^{|\alpha^{(0)}| - N}. \quad (6)$$

Consider a cycle C in E including kr_1 edges of which r_2 are negative. As in the case $k = 1$, if C is an e-cycle, then $\text{sign}(C) = (-1)^{r_1}$, while if C is an o-cycle, then $\text{sign}(C) = (-1)^{r_1+1}$. Decompose E into disjoint cycles, which comprise N_e e-cycles, and N_o o-cycles. Let $k\theta_0$ be the total number of edges in o-cycles and $k\theta_e$ the total number of edges in e-cycles, so that $\theta_o + \theta_e = |\alpha^{(0)}|$ (since there are $k|\alpha^{(0)}|$ edges in E). Taking the product of signs of edges in E over e-cycles and o-cycles separately gives

$$\prod_{j=0}^{k-1} \text{sign} \left(A_{\alpha^{(j)}, \beta^{(j+1)}}^{(j)} \right) = (-1)^{\theta_o} (-1)^{\theta_e + N_o} = (-1)^{|\alpha^{(0)}| + N_o}. \quad (7)$$

Substituting Eqs. 6 and 7 into Eq. 5 gives:

$$\text{sign}(T) = (-1)^{|\alpha^{(0)}| - N} (-1)^{|\alpha^{(0)}| + N_o} = (-1)^{N_e}. \quad (8)$$

Note that this is just Eq. 4 again. As in the case $k = 1$, if G is e-cycle-free, then $\text{sign}(T) = 1$, and since T is an arbitrary nonzero term in $A[\alpha^{(0)}]$, $A[\alpha^{(0)}] \geq 0$. Since $\alpha^{(0)}$ is an arbitrary nonempty subset of $\{1, \dots, n_0\}$, A is a P_0 -matrix. \square

We have the following corollary to Theorem 1:

Corollary 4. *Consider a square matrix $A = A^{(0)}A^{(1)} \dots A^{(k-1)}$ such that $G = G_{A^{(0)}A^{(1)} \dots A^{(k-1)}}$ is e-cycle-free. Then matrices in $\mathcal{Q}(A^{(0)})\mathcal{Q}(A^{(1)}) \dots \mathcal{Q}(A^{(k-1)})$ are all P_0 -matrices.*

PROOF. By definition, any matrix $B \in \mathcal{Q}(A^{(0)})\mathcal{Q}(A^{(1)}) \dots \mathcal{Q}(A^{(k-1)})$ can be written $B = B^{(0)}B^{(1)} \dots B^{(k-1)}$, where $B^{(j)} \in \mathcal{Q}(A^{(j)})$. But

$$G_{B^{(0)}B^{(1)} \dots B^{(k-1)}} \cong G_{A^{(0)}A^{(1)} \dots A^{(k-1)}},$$

so, by Theorem 1, B is a P_0 -matrix. \square

Remark. For each $r = 1, \dots, k-1$ the matrix product

$$A^{(r)}A^{(r+1)} \dots A^{(k-1)}A^{(0)} \dots A^{(r-1)}$$

gives rise to a graph isomorphic to $G = G_{A^{(0)}A^{(1)} \dots A^{(k-1)}}$. Thus, in fact, if G is e-cycle-free, then all matrices in

$$\mathcal{Q}(A^{(r)})\mathcal{Q}(A^{(r+1)}) \dots \mathcal{Q}(A^{(k-1)})\mathcal{Q}(A^{(0)}) \dots \mathcal{Q}(A^{(r-1)})$$

are P_0 -matrices.

5. Converse results and proof of Theorem 2

A variety of converse results are possible, that is, results which guarantee that if a signed $(k, \{1\})$ -BC graph contains e-cycles, then there exist matrices in some set which fail to be P_0 -matrices. The most useful formulations depend on the application. Lemma 5 is the basic result from which such results follow:

Lemma 5. *Consider a graph $G = G_{A^{(0)}A^{(1)}\dots A^{(k-1)}}$ such that all edges in G lie on a single e-cycle C of length kr . Then $A = A^{(0)}A^{(1)}\dots A^{(k-1)}$ is not a P_0 -matrix.*

PROOF. Define the sets $(\alpha^{(0)}, \dots, \alpha^{(k-1)})$ by the stipulation that $s \in \alpha^{(i)}$ if and only if V_i^s lies on C . Clearly $|\alpha^{(i)}| = r$ for each i . Each vertex in $V_j^{\alpha^{(j)}}$ lies on C and hence has exactly two edges, one incoming, and one outgoing, incident on it. So it is possible to define bijections $\phi_j : V_j^{\alpha^{(j)}} \rightarrow V_{j+1}^{\alpha^{(j+1)}}$ as follows: given vertices $v \in V_j^{\alpha^{(j)}}$ and $\tilde{v} \in V_{j+1}^{\alpha^{(j+1)}}$, $\phi_j(v) = \tilde{v}$ if there is a directed edge (v, \tilde{v}) in E . Each ϕ_j induces a permutation $\beta^{(j+1)} : V_{j+1}^{\alpha^{(j+1)}} \rightarrow V_{j+1}^{\alpha^{(j+1)}}$ defined by $\beta^{(j+1)} = \phi_j \circ \iota_j^{-1} \circ \iota_{j+1}$ (see Figure 2).

Consider the minor $A[\alpha^{(0)}]$. There is a nonzero term in this minor

$$T = \prod_{j=0}^{k-1} T_j = \prod_{j=0}^{k-1} \sigma(\beta^{(j+1)}) A_{\alpha^{(j)}, \beta^{(j+1)}}^{(j)}.$$

Moreover T is the unique nonzero term in $A[\alpha^{(0)}]$: any other nonzero term would imply the existence of an index j and a permutation $\delta : V_{j+1}^{\alpha^{(j+1)}} \rightarrow V_{j+1}^{\alpha^{(j+1)}}$, $\delta \neq \beta^{(j+1)}$, such that $A_{\alpha^{(j)}, \delta}^{(j)} \neq 0$. Letting s be an index such that $\delta_s \neq \beta_s^{(j+1)}$, $A_{\alpha_s^{(j)}, \delta_s}^{(j)}$ must then be nonzero, implying the existence of an edge $(V_j^{\alpha_s^{(j)}}, V_{j+1}^{\delta_s})$ in G which does not lie in C . But by assumption C contains all edges in G .

By Eq. 8, $\text{sign}(T) = (-1)^{N_e}$, where N_e is the number of e-cycles in the subgraph associated with T . Since this subgraph is precisely C , $N_e = 1$ and $\text{sign}(T) = -1$. Thus $A[\alpha^{(0)}] < 0$ and A fails to be a P_0 -matrix. \square

Corollary 6 illustrates an application of Lemma 5:

Corollary 6. *Suppose a graph $G_{A^{(0)}A^{(1)}\dots A^{(k-1)}}$ contains an e-cycle C . Then there are matrices in $\mathcal{X} \equiv \mathcal{Q}(A^{(0)})\mathcal{Q}(A^{(1)})\dots\mathcal{Q}(A^{(k-1)})$ which are not P_0 -matrices.*

PROOF. Each edge in C corresponds to an entry in one of the matrices $A^{(j)}$. For each j define $\tilde{A}^{(j)}$ to be the matrix $A^{(j)}$ with all entries *not* corresponding to edges in C set to be zero. Then the matrix factorisation $\tilde{A} = \tilde{A}^{(0)}\tilde{A}^{(1)}\dots\tilde{A}^{(k-1)}$ gives rise to a graph which consists solely of the e-cycle C , and hence, by Lemma 5, \tilde{A} fails to be a P_0 -matrix. But $\tilde{A} \in \text{cl}(\mathcal{X})$ (that is the closure of \mathcal{X}), and since the set of P_0 -matrices is closed, there are matrices in \mathcal{X} which fail to be P_0 . \square

PROOF OF THEOREM 2. This is immediate: Theorem 2 is simply a combination of Corollaries 4 and 6. \square

6. Notes and conclusions

Although the special case $k = 1$ was treated for completeness, the result can easily be inferred from previous work. For $k = 1$, Theorem 2 states that given a square matrix A , all matrices in $\mathcal{Q}(A)$ are P_0 -matrices if and only if the (unique) signed $(1, \{1\})$ -BC digraph G_A associated with A is e-cycle-free. However G_A is closely related to an object often called the interaction graph or I-graph in the literature. In fact the I-graph associated with A is just G_{A^T} . Results in [9, 12] showed that G_{-A^T} lacks positive cycles if and only if all matrices in $\mathcal{Q}(A)$ are P_0 -matrices. Trivially, G_{-A^T} lacks positive cycles if and only if G_{-A} lacks positive cycles. The definitions imply that e-cycles (resp. o-cycles) in G_A are in one-to-one correspondence with positive cycles (resp. negative cycles) in G_{-A} . So G_{-A} lacks positive cycles if and only if G_A is e-cycle-free. Together these observations imply that matrices $\mathcal{Q}(A)$ are all P_0 -matrices if and only if G_A is e-cycle-free.

The case $k = 2$ has also effectively been treated previously in [11, 12], where the associated graphs were termed “DSR graphs”. The main differences between the definition of a DSR graph in [12], and a signed $(2, \{1\})$ -BC digraph here, are (i) directions on all edges are reversed, (ii) here, edge-labels have been ignored, while some computations on DSR graphs in [11, 12] involved edge-labels, and (iii) in the construction of the DSR graph a pair of identically signed edges (\tilde{v}, v) and (v, \tilde{v}) are replaced with a single undirected edge, with a view to removing o-cycles of length 2 from the graph, and thus simplifying computation. This process neither creates nor destroys e-cycles, and so does not change the key fact that an absence of e-cycles implies that associated matrices are P_0 -matrices.

The treatment in [11, 12] also suggests that extensions obtaining sharper results by introducing edge-labelling and more complex computations on the graphs are possible. The most useful forms that such extensions might take depend on the applications in question. These directions will be treated in future work.

Appendix. Definitions

Matrices

- **P -matrix**: a square matrix all of whose principal minors are positive.
- **P_0 -matrix**: a square matrix all of whose principal minors are nonnegative.
- **Qualitative class of a matrix**. Given a matrix M , the qualitative class of M , $\mathcal{Q}(M)$, is the set of matrices of the same dimensions as M , and with the same sign pattern as M .

Digraphs

- **Digraph:** a pair $G = (V, E)$, where V is a set of vertices, and E is a set of directed edges, i.e., each element of E is an ordered pair of vertices.
- **Signed digraph:** a pair (G, σ) , where G is a digraph, and σ is a sign function on the edge set E of G , i.e., $\sigma : E \rightarrow \{-1, +1\}$.
- **Cycle (in a digraph):** a simple directed path from a vertex to itself.
- **Signed (k, S) -block circulant digraph.**
 - i. $k = 1$: any signed digraph.
 - ii. $k \geq 2$: a signed digraph whose vertices can be partitioned into k sets $\{V_0, V_1, \dots, V_{k-1}\}$, such that if $(j - i) \bmod k \notin S$, then there is no (directed) edge from a vertex in V_i to a vertex in V_j .
- **e-cycle/o-cycle** in a signed $(k, \{1\})$ -block circulant digraph. A cycle with kr_1 edges, of which r_2 have negative sign is an e-cycle if $(-1)^{r_1+r_2} = 1$ and an o-cycle if $(-1)^{r_1+r_2} = -1$.

Permutations

- **Permutation.** Given an ordered set $\alpha = [\alpha_1, \dots, \alpha_n]$, a permutation of α is used variously to refer to a bijection $\beta : \alpha \rightarrow \alpha$, or to the (ordered) image of this bijection $[\beta(\alpha_1), \dots, \beta(\alpha_n)]$.
- **Sign of a permutation.** A permutation α is even (resp. odd) if it can be decomposed as an even (resp. odd) number of transpositions. Its sign $\sigma(\alpha)$ is defined by: $\sigma(\alpha) = +1$ if α is an even permutation, and $\sigma(\alpha) = -1$ if α is odd.

- [1] D. Hershkowitz and N. Keller. Positivity of principal minors, sign symmetry and stability. *Linear Algebra Appl.*, 364:105–124, 2003.
- [2] K.G. Murty. *Linear Complementarity, Linear and Nonlinear Programming*. Heldermann Verlag, Berlin, 1988.
- [3] R.W. Cottle, J.-S. Pang, and R.E. Stone. *The Linear Complementarity Problem*. Academic Press, New York, 1992.
- [4] J. Hofbauer and K. Sigmund. *Evolutionary games and population dynamics*. Cambridge University Press, 1998.
- [5] D. Gale and H. Nikaido. The Jacobian matrix and global univalence of mappings. *Math. Ann.*, 159:81–93, 1965.
- [6] T. Parthasarathy. *On global univalence theorems*, volume 977 of *Lecture Notes in Mathematics*. Springer-Verlag, 1983.
- [7] M. Banaji, P. Donnell, and S. Baigent. P matrix properties, injectivity and stability in chemical reaction systems. *SIAM J. Appl. Math.*, 67(6):1523–1547, 2007.

- [8] P. Donnell, M. Banaji, and S. Baigent. Stability in generic mitochondrial models. *J. Math. Chem.*, 46(2):322–339, 2009.
- [9] C. Soulé. Graphic requirements for multistationarity. *Complexus*, 1:123–133, 2003.
- [10] M. Banaji and G. Craciun. Graph-theoretic criteria for injectivity and unique equilibria in general chemical reaction systems. *Adv. in Appl. Math.*, 44:168–184, 2010.
- [11] M. Banaji and G. Craciun. Graph-theoretic approaches to injectivity and multiple equilibria in systems of interacting elements. *Commun. Math. Sci.*, 7(4):867–900, 2009.
- [12] M. Banaji. Graph-theoretic conditions for injectivity of functions on rectangular domains. *J. Math. Anal. Appl.*, 370:302–311, 2010.
- [13] R. A. Brualdi and B. L. Shader. *Matrices of sign-solvable linear systems*. Number 116 in Cambridge tracts in mathematics. Cambridge University Press, 1995.
- [14] B. Alspach and T. D. Parsons. Isomorphism of circulant graphs and digraphs. *Discrete Math.*, 25(2):97–108, 1979.
- [15] F. Boesch and R. Tindell. Circulants and their connectivities. *J. Graph Theory*, 8:487–499, 1984.
- [16] F. R. Gantmacher. *The theory of matrices*. Chelsea, 1959.