

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Bishop, Chris, Turner, Anthony N., Jarvis, Paul, Chavda, Shyam and Read, Paul (2017)
Considerations for selecting field-based strength and power fitness tests to measure
asymmetries. *Journal of Strength and Conditioning Research*, 31 (9). pp. 2635-2644. ISSN
1064-8011

Final accepted version (with author's formatting)

This version is available at: <http://eprints.mdx.ac.uk/22024/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

1 **CONSIDERATIONS FOR SELECTING FIELD-BASED STRENGTH**
2 **AND POWER FITNESS TESTS TO MEASURE ASYMMETRIES**

3
4 **AUTHORS:**

5 Chris Bishop¹ (MSc), Anthony Turner¹ (PhD, CSCS*D), Paul Jarvis¹ (MSc), Shyam
6 Chavda¹ (MSc, CSCS) and Paul Read^{2,3} (PhD, CSCS*D)

7
8 **AFFILIATIONS:**

9 1. School of Science and Technology, London Sports Institute, Middlesex University,
10 London, UK

11 2. Athlete Health and Performance Research Centre, Aspetar Orthopaedic and Sports
12 Medicine Hospital, Doha, Qatar

13 3. Youth Physical Development Centre, School of Sport, Cardiff Metropolitan University,
14 Cardiff, UK

15
16 **CORRESPONDENCE:**

17 **Name:** Chris Bishop

18 **Address:** 10 Picketts, Welwyn Garden City, Hertfordshire, AL8 7HJ, UK

19 **Email:** C.Bishop@mdx.ac.uk

1 **Abstract**

2 The prevalence of lower limb asymmetries has been reported in numerous studies; however,
3 methodological differences exist in the way they can be detected. Strength and jumping-
4 based tasks have been most commonly used to examine these differences across both athlete
5 and non-athlete populations. The aim of this review was to critically analyze the utility of
6 strength and jumping tests that are frequently used to measure asymmetry. Reliability,
7 validity, and considerations for assessment are examined to enhance test accuracy and
8 effectiveness in the quantification of asymmetries during strength and jumping-based tasks.
9 Medline and SPORT Discus databases were used with specific search terms to identify
10 relevant articles in both athlete and non-athlete populations. The findings of the current
11 review indicate that assessing inter-limb differences during strength and jumping-based tasks
12 may result in different levels of asymmetry; thus, inter-limb differences appear to be task-
13 dependent. Consequently, quantification during both types of assessment is warranted and a
14 selection of tests has been suggested to measure asymmetries in both strength and jumping-
15 based tasks.

16

17 **Key Words:** Asymmetries, Reliability, Strength, Jumping

18

19

20

21

22

1 **Introduction**

2 The concept of asymmetries has attracted much interest in strength and conditioning (S&C)
3 in recent years. Multiple studies have reported the prevalence of asymmetries during a variety
4 of jumping (7,23,32,65) and strength-based assessments (4,24,47,56,58); however, a critical
5 analysis of their utility for measuring inter-limb differences and clear guidelines for
6 implementation are sparse. Within the available literature, methodological differences exist
7 with regards to the type of test used and their administration procedures. The
8 countermovement jump (CMJ) and single leg CMJ (SLCMJ) have been most commonly used
9 (7,13,36,43,47,63), most likely due to their ease of application. However, solely measuring
10 lower body jumping-based tasks in a vertical direction will reduce the ecological validity for
11 a range of sports which require movements in multiple planes of motion and a range of
12 physical qualities. Previous data also indicate that measures of strength, such as the back
13 squat (22,47,58), isometric squat or mid-thigh pull (IMTP) (4,25,30,65), and isokinetic knee
14 flexion or extension (15,18,56,59) have shown adequate sensitivity to identify between-limb
15 differences. Furthermore, any highlighted inter-limb differences in strength and jumping
16 tasks have shown decrements in physical performance (4,7,69), increased injury risk (35),
17 and reduced performance in sport-specific tasks (26). Therefore, to provide an accurate
18 profile to screen athletes for the presence of asymmetry, a battery of tests may be required
19 due to the potential for task sensitivity across a range of physical competencies.

20 When calculating asymmetries, a variety of approaches have been used which define limb
21 differences in terms of dominance, strength, preference, or simply a right or left distinction
22 (10). For example, studies pertaining to soccer frequently define the dominant limb as the
23 favoured 'kicking leg' (15, 56), which seems valid considering the nature of such a task.
24 However, recent research has highlighted poor levels of agreement (40%) between perceived
25 limb dominance and the highest score attained (23). In addition, Zifchock et al. (70)

1 suggested that numerous asymmetry equations emphasise the use of a ‘reference value’ (such
2 as the dominant limb or highest score); however, clarity is sometimes lacking as to why one
3 limb is chosen over the other. Therefore, precision on defining limb dominance is critical and
4 must retain specificity to the task in question.

5 A combination of factors exist that should be considered prior to the selection of appropriate
6 tests to measure asymmetry. These include test reliability to ensure there is adequate
7 precision in the outcome measures used to enable practitioners to accurately quantify inter-
8 limb differences. Associations with reductions in performance or heightened injury risk
9 should also be considered in determining the usefulness of a test. Finally, the requirements of
10 the athlete must be considered within the context of their sport. For example, ski athletes
11 perform their sport bilaterally and it may be logical to choose bilateral tests when quantifying
12 asymmetries in strength and jumping tasks (38). However, team sports such as soccer and
13 rugby hold a greater degree of unpredictability in an athlete’s movement patterns; thus,
14 unilateral testing or a combination of both may be most applicable. Additional reasons such
15 as experience of the tester, ease of testing equipment and cost effectiveness should also be
16 considered and will be discussed later in this review.

17 The aim of this review was to provide an overview of the current literature to critically
18 examine which tests could be considered when quantifying asymmetries in strength and
19 jumping-based tasks. Sub-headings of ‘reliability’, ‘associations with athletic performance’
20 and ‘athlete requirements’ have been provided to promote a clear rationale for suggestions
21 made. Additional sections on how to interpret change (in asymmetry values) and practical
22 considerations for testing will help to provide the reader with context on subsequent data
23 analysis and expectations during test protocols. Finally, a test battery has been proposed for
24 the assessment of asymmetries in strength and jumping-based tasks.

1

2 **Methods**

3 Empirical research studies and review journal articles were retrieved from electronic searches
4 of Medline and SPORT Discus databases. The search strategy chose to combine specific
5 terms with the word 'asymmetries' to ensure relevant articles were extracted. The search
6 terms included: 'asymmetries and strength', 'asymmetries and jumping', 'asymmetries and
7 performance', 'reliability of strength tests' and 'reliability of jumping tests'. Articles were
8 deemed relevant after scanning the title and abstract and where subsequent access to the full
9 text was available from the relevant publishers. The reference lists of each study were also
10 checked to ensure no further articles were omitted from the search process. All searches were
11 conducted between the 1st November and 30th November, 2016.

12

13 **Strength Tests**

14 Reliability

15 The back squat has been used in a number of studies to examine the prevalence of force
16 production asymmetries (22,31,47,58). Considering the bilateral nature of the exercise, any
17 data pertaining to right and left differences would likely require the use of twin force plates.
18 Strong reliability has been shown when testing bilateral 1RM loads for both novice and
19 experienced lifters (ICC [intraclass correlation coefficient] = 0.94) (53) and healthy male
20 (ICC = 0.99, CV = 0.35%) and female subjects (ICC = 0.97, CV = 0.53%) (60). Previously, a
21 CV target of < 10% has been deemed acceptable (14); thus, it would appear that the 1RM
22 back squat is a reliable method for testing strength across genders and experienced or
23 inexperienced lifters. However, to the authors' knowledge, only two studies have investigated

1 data pertaining to vertical ground reaction force (vGRF) asymmetries during the back squat.
2 Newton et al. (47) used 14 NCAA softball players to perform three back squats at 80% 1RM
3 and reported average GRF asymmetries of 6.02%. Hodges et al. (31) examined vGRF
4 asymmetry during the first and last two repetitions in each set of a training session that was
5 comprised of 5 sets of 8 repetitions at 90% of their 8RM. Mean inter-limb differences (across
6 all sets) were reported to be 4.3% for the first two repetitions and 3.6% for the final two
7 repetitions. The results from these two studies indicate that vGRF asymmetries are low
8 during the back squat for college and healthy adult populations; although further research is
9 required to examine the reliability of inter-limb differences during this test.

10 The isometric squat or IMTP have also been used to measure asymmetry in the available
11 research (4,5,6,19,25). Bilateral versions of these tests also require the use of twin force
12 plates with peak vGRF most commonly reported (4,6,25). In addition, other variables that
13 may be of interest to S&C coaches such as rate of force development (RFD) and impulse can
14 also be calculated (19,25,40). Due to the restricted timeframe within sporting movements that
15 athletes have to produce force (1), these physical characteristics can be considered an
16 important diagnostic; however, the reliability of measurement may be questionable. Hart et al.
17 (25) measured the reliability of peak force, mean force and RFD during bilateral and
18 unilateral isometric squats. Results showed that rank-order repeatability via the ICC and
19 within-subject variation via the CV for peak and mean force were acceptable based on
20 previous guidelines (14). RFD was more variable (see Table 1), although it should be noted
21 that this was on the non-dominant limb only. Furthermore, the subjects used in this study
22 were not of a specific sporting background and as such may produce more variation in their
23 results due to a possible lack of familiarity with testing protocols, which has been seen
24 elsewhere (57).

1 *** *INSERT TABLE 1 ABOUT HERE* ***

2

3 For the IMTP, Dos Santos et al. (19) investigated the prevalence of strength asymmetries
4 between professional rugby league and collegiate athletes. All subjects performed three
5 bilateral and unilateral trials on each limb with peak force and impulse reported. Results
6 showed strong reliability for unilateral peak force (ICC = 0.94; CV = 4.7-5.0%), but more
7 variability for impulse (ICC = 0.82-0.88; CV = 9.3-11.6%). Significant differences ($p < 0.05$)
8 between dominant and non-dominant limbs for both groups of athletes were reported,
9 suggesting that the unilateral IMTP was a reliable method for determining strength
10 asymmetries across athletes of different levels (19). In addition, reliability data has also been
11 reported for both males ($n = 31$) and females ($n = 32$) during the IMTP. Bailey et al. (5)
12 reported an ICC range of 0.68-0.98 for multiple variables including peak force, impulse at
13 different time points, and RFD although individual ICC values were not specified for the
14 tested metrics. The standard error of measurement (SEM), which is an indication of a score's
15 accuracy (68), was also reported and the highest variability was noted for impulse at 50 m/s.
16 Although individual ICC's were not reported, the SEM is a measure of absolute reliability
17 and thus, is arguably a more important measure. With that in mind, lower levels of reliability
18 for impulse are in agreement with the findings of Dos Santos et al. (19). Furthermore, the
19 sample was divided into stronger and weaker sub-groups with SEM values significantly
20 different ($p < 0.05$) between groups for peak force (0.07 vs. 0.13 s) and RFD (0.45 vs. 0.70
21 n/s^{-1}). The authors stated that strength may be a more influential factor than sex when
22 calculating asymmetries during the IMTP due to the increased variability and inter-limb
23 differences seen in the weaker group (5).

1 Isokinetic dynamometry is another alternative for practitioners who wish to measure both
2 inter- and intra-limb strength asymmetries in isolated joint actions (such as knee flexion or
3 extension). Research is available to analyse the presence of asymmetries in different
4 populations ranging from collegiate (36,41,47) to professional athletes (15,55,59); however,
5 none of these studies included data examining the reliability of inter-limb differences.
6 Impellizzeri et al. (34) reported the reliability of isokinetic strength imbalance ratios (between
7 hamstrings and quadriceps) showing weak to strong reliability (ICC range = 0.34-0.87).
8 Furthermore, the SEM ranged from 3.2-8.7% for strength imbalance ratios and 4.3-7.7% for
9 peak torque measurements. Similar reliability values have also been reported for intra-limb
10 isokinetic knee flexion and extension measurements in recreational athletes (3). Concentric
11 and eccentric actions were recorded at 60, 180 and 240°·sec⁻¹ with reliability assessed via the
12 SEM and ICC. All measures showed moderate reliability with the percentage error reaching
13 as high as 20% and ICC's > 0.7 (3).

14

15 Effects on Athletic Performance

16 There is currently a paucity of literature to examine the relationship between asymmetry
17 during the aforementioned strength tests and athletic performance. Bailey et al. (4) reported
18 mean asymmetries during the IMTP of 6.6%, and moderate negative correlations between the
19 peak force symmetry index and jump height ($r = -0.39$ to -0.52) and peak power ($r = -0.28$ to
20 -0.43) across loaded (20 kg) and unloaded jumps. Furthermore, Rannama et al. (51) showed
21 that peak torques asymmetries of the knee extensors (measured at 180°·sec⁻¹) were negatively
22 correlated ($r = -0.50$; $p < 0.05$) with power during a 5-second maximal effort cycling test.
23 Strength asymmetries have also been shown to have a detrimental effect on the performance
24 of sport-specific actions. Hart et al. (26) used a unilateral isometric squat to determine inter-

1 limb strength differences in Australian football players. Higher asymmetries (8%) were
2 negatively associated with kicking accuracy compared to the accurate players who only
3 exhibited 1% imbalances in strength. These data indicate that larger strength asymmetries
4 may have a negative impact on performance; however, caution should be applied due to high
5 amount of variance remaining unexplained by these relationships.

6

7 *Athlete Requirements*

8 When selecting appropriate tests to measure asymmetry, practitioners should consider their
9 ecological validity. For example, bilateral assessments may be more suitable for a powerlifter,
10 to ensure task specificity is being adhered to. Conversely, team sport athletes are required to
11 undertake multiple unilateral sporting actions such as running and changing direction;
12 therefore, it seems logical to suggest some form of unilateral strength testing when
13 calculating asymmetries.

14 The type of muscle actions and speeds of movement involved in the sport are also a
15 consideration in test selection. Isokinetic testing has the advantage of measuring asymmetries
16 across a range of muscle actions (concentric and eccentric) and speeds unilaterally,
17 potentially providing a more complete picture of strength asymmetries. However, strength
18 during single joint actions is not fully representative of compound movement patterns (9,36),
19 which are more characteristic of the actions required during the execution of the majority of
20 sporting tasks. Furthermore, isokinetic dynamometry testing requires a laboratory and
21 expensive equipment which may not be practically viable for many athletes or teams. Until
22 recently, it could have been argued that this notion held true for the use of force plates;
23 however, more recently affordable (and portable) versions are now available increasing their
24 utility for field testing large numbers of athletes.

1

2 **Jump Tests**

3 Reliability

4 When determining asymmetries in jump tests, a variety of bilateral and unilateral tests have
5 been frequently used (7,13,35,36,41,49,52,54,69), most likely because of their ease of
6 implementation. Although inter-limb differences can be calculated with only one force plate,
7 as per the methods of Impellizzeri et al (35), large movement variability in vertical jumping
8 has been noted (8,38), perhaps suggesting that asymmetries should be determined within the
9 same repetition if quantified bilaterally. Alternatively, asymmetries can also be measured via
10 single leg jumping tasks through the use of a jump mat and assessment of flight time and
11 ground contact duration. Whilst not a common topic, Benjanuvatra et al. (8) aimed to
12 differentiate between the bilateral CMJ and SLCMJ for assessing asymmetries in impulse and
13 vGRF. The authors suggested using the SLCMJ over the bilateral CMJ when quantifying
14 asymmetries because it places a greater emphasis on force production from one limb and
15 reduces the athlete's base of support presenting a challenge that is more representative of the
16 actions performed for most sports. Furthermore, multiple sporting actions such as jumping
17 and sprinting occur unilaterally; thus, the notion of specificity is kept to the sporting task if
18 asymmetries are tested for unilaterally. Therefore, single leg tasks may provide a more
19 accurate reflection of true inter-limb asymmetries for team sport athletes in particular.
20 However, it should be noted that task specificity may ultimately dictate which jump test is
21 chosen.

22 Meylan et al. (46) reported strong reliability for measures of jump height and distance during
23 the SLCMJ and lateral jumps. ICC's ranged from 0.91-0.98 across both genders in healthy
24 adults. Furthermore, CV ranges fell between 2.7-7.2%, suggesting that multi-directional,

1 unilateral jumps are a reliable method for assessing between-limb differences. Strong
2 reliability has also been noted in youth athletes for measures of peak force and power during
3 the SLCMJ (13), with ICC's ranging from 0.88-0.97. Consequently, unilateral vertical jump
4 assessments appear to be reliable tests across adult and youth populations.

5 The reliability of various single leg hop tests has also been measured within previous research
6 (12,52,55). Common variations include the single leg hop (for distance), triple hop, 6m timed
7 hop, and crossover hop (Figure 1). The single leg hop would appear to be the most reliable of
8 these four tests with ICC's ranging from 0.92-0.96 and SEM's of 4.56-4.61cm, with more
9 variability present in the 6m timed hop (ICC = 0.66-0.92) (12,52,55). Despite their
10 similarities, it has been suggested that more than one hop test should be considered when
11 quantifying asymmetries (48) because of the different demands they each pose. Considering
12 the previously reported strong reliability of the triple hop test (ICC = 0.88-0.97), and notably
13 lower SEM values when compared to the crossover hop (11.17 vs. 17.74 cm) (52,55), the
14 rebound nature of the task may provide a more ecologically valid representation of unilateral
15 tasks for athletes in running and jumping based sports. However, it must be acknowledged
16 that this test likely places a greater physical demand on athletes and should be used with
17 caution if plyometric training experience is low. Furthermore, it is plausible that a low
18 training age for this physical quality may negatively affect the reliability of the test, rendering
19 it unsuitable; although further research is again warranted to substantiate this theory.

20

21 *** *INSERT FIGURE 1 ABOUT HERE* ***

22

1 The reliability of various measures collected during drop jumping tasks has also been
2 reported in the available research (2,17,21,67), although it would appear only recently that
3 this exercise has been used to report asymmetries. Maloney et al. (44) showed bilateral drop
4 jump asymmetries as high as 59.7% for leg stiffness, whilst within-session reliability (CV)
5 was 5% for vGRF. However, CV's were noticeably higher for negative centre of mass
6 displacement and vertical stiffness (12 and 13% respectively), although this may have been
7 attributed to the sample not being an athletic population and therefore test familiarity must be
8 questioned. Although not used for asymmetry detection, test-retest reliability (using the ICC)
9 has previously been reported in the drop jump for measures of peak and mean force (0.86-
10 0.98), jump height (0.99), and ground contact times (0.98) (17,21), indicating strong rank-
11 order repeatability. However, further research is warranted to examine the reliability of these
12 variables with respect to asymmetry.

13 While the majority of the available literature pertaining to the reliability of drop jumping
14 tasks is focused on bilateral variations, the unilateral drop jump has also recently been
15 examined (44). The authors reported similar levels of asymmetry as the bilateral test (~55%)
16 and showed within-session CV's of 2% for vGRF, indicating small variability between trials
17 in a non-athletic population. Stalboom et al. (62) investigated the reliability of impulse, mean
18 and peak force during single leg drop jumping and found ICC's ranged from 0.74-0.96 and
19 all CV's < 10%. Although both studies indicate acceptable levels of reliability, procedures
20 were conducted from 18 and 20cm boxes respectively. Bilateral drop jump measures are
21 frequently conducted from a height of 30cm (21,39,42), but the increased physical demand
22 associated with a unilateral version would suggest that lower heights may be more
23 appropriate. This is supported by Maloney et al. (44) who described how the required short
24 ground contact times could not be maintained when dropping from heights of 30 and 45cm.

1 Effects on Athletic Performance

2 Lockie et al. (43) investigated asymmetries in different unilateral jump tasks and their
3 relationships with performance tests. Between limb-differences of 10.4%, 3.3% and 5.1%
4 were shown for CMJ height, broad jump and lateral jump distances respectively. No
5 significant correlations were found between any of the asymmetry values and sprint (5, 10,
6 20m) or change of direction speed (CODS) tests (left and right 505 and modified t-test).
7 These data indicate that asymmetries $\leq 10\%$ do not negatively affect sprint of COD
8 performance. This is supported by Hoffman et al. (32), with SLCMJ asymmetries of 9.7%
9 showing no significant differences in time when performing the 3-cone drill to the dominant
10 or non-dominant side. Significant weak correlations were reported for the SLCMJ non-
11 dominant limb and the 3-cone drill for both dominant ($r = -0.36; p < .05$) and non-dominant
12 ($r = -0.37; p < .05$) directions; yet no significance was found when compared with the
13 dominant limb of the SLCMJ. This suggests that asymmetries may be task dependent and
14 highlights the multi-faceted nature of CODS that requires multiple factors to determine its
15 success (61).

16 In contrast, inter-limb differences measured during jump tests have been shown to
17 detrimentally impact performance (28). Significant correlations were reported between
18 unilateral jump test asymmetries (vertically, horizontally and laterally) and measures of mean
19 ($r = 0.40-0.43$) and peak force ($r = 0.38-0.48$) during a reactive agility task. Furthermore, the
20 dominant limb was significantly faster for agility push-off time (1.48 vs. 1.56 sec; $p < .001$)
21 compared to the non-dominant side (28). Bell et al. (7) investigated how lean muscle mass
22 asymmetries affected CMJ performance. Results portrayed that thigh and shank lean mass
23 asymmetry accounted for 20% of the variance in force asymmetry. Further to this, lean mass
24 asymmetry of the pelvis, thigh, and shank accounted for 25% of power asymmetries, both
25 during the CMJ. Additionally, asymmetries in power $> 10\%$ during the CMJ resulted in

1 decreased jump height of 9cm (effect size = $d > 0.8$), which represents a considerable
2 reduction in jump performance. Maloney et al. (45) attempted to determine the relationship
3 that asymmetries (during single leg drop jumps) had with a 90° cutting task (performed on a
4 force plate). The sample was subsequently divided into slow and fast groups ($n = 9$ per
5 group), with mean vertical stiffness and jump height asymmetry explaining 63% of the
6 cutting task variance ($r^2 = 0.63$; $p = .001$). Furthermore, faster athletes portrayed significantly
7 lower asymmetries for jump height ($p = .026$), perhaps indicating that minimizing differences
8 in reactive strength could be advantageous to cutting performance also. These results support
9 the notion that it may be advantageous to minimize inter-limb differences when aiming to
10 optimize agility and jumping performance.

11 Finally, the majority of research pertaining to asymmetries in jump tests often report data
12 relating to performance outcomes (such as jump height or distance) or propulsive forces.
13 Intra-limb differences in eccentric forces or mechanics (such as landing in this instance) are
14 seldom reported and should also be considered to build the asymmetry picture further. Pappas
15 and Carpes, (50) showed that knee valgus was significantly greater when jumping forward
16 (20cm) compared to a drop land (from a 40cm box), highlighting the notion of task-
17 specificity for landing kinematics as well. Whilst minimal research pertaining to asymmetries
18 during landing patterns currently exists, the relationship between eccentric forces and injury
19 risk is recognised (15,36); thus, further research in this area for inter-limb differences is
20 warranted.

21

22 *Athlete Requirements*

23 Similar to testing for asymmetries in strength, both the sporting needs analysis and individual
24 athlete requirements should dictate which tests are chosen. However, the concept of

1 movement variability must be considered during jump testing and will have considerable
2 impact on how asymmetries are interpreted. Exell et al. (20) outlined how an asymmetry can
3 only be considered “real” if the inter-limb differences are greater than the variability within
4 an athlete’s movement. This concept was originally investigated during running; however,
5 Jordan et al. (38) noted a shift in how asymmetries presented themselves during both the
6 eccentric and concentric phases of a CMJ, supporting this notion of movement variability in
7 jumping. With that in mind, it is essential that coaches understand how to calculate
8 movement variability. A well organised protocol will often assume three trials of a test (66),
9 and the CV allows practitioners to quantify the variation between trials for said test
10 (expressed as a percentage). With asymmetries also being reported as a percentage, this
11 allows a comparison of asymmetry score compared to variability score, and ultimately will
12 help to determine if an asymmetry is real. Therefore, for athletes that require multiple
13 movement patterns in different planes of motion (29,30), it is suggested that practitioners
14 consider initial testing both bilaterally and unilaterally. Subsequent data analysis such as the
15 CV (which also serves as an indicator of variability) may provide an impression as to whether
16 certain tests are deemed too unreliable for their athlete population, which will help to
17 streamline future asymmetry testing. As outlined by Cormack et al. (14), practitioners should
18 consider $< 10\%$ as a target threshold when determining the reliability of a test and this figure
19 can then serve as a comparison to the subsequent asymmetry value.

20

21 **Interpreting Asymmetry Scores**

22 Determining critical thresholds for asymmetry that are linked to reductions in performance or
23 heightened injury risk provides S&C coaches with useful data to design targeted training
24 interventions for athletes. The available body of literature suggests that asymmetries are task-

1 specific; meaning that practitioners should not expect to see the same inter-limb differences
2 across different tests for the same physical quality. This is supported by Jones and
3 Bampouras, (36) who reported that asymmetries varied across tasks with differences of 4.47%
4 and 12.43% for jump and strength tests respectively. Furthermore, Schiltz et al. (59) reported
5 strength and power asymmetries of 6.5 and 12% in professional basketball players during
6 isokinetic and drop jump testing respectively; justifying undertaking tests across both
7 physical competencies.

8 Where strength asymmetries are concerned, Bailey et al. (4) reported negative associations
9 with jump performance when strength differences of 6.6% were seen from the IMTP. Hart et
10 al. (26) noted significantly reduced performance in kicking accuracy with asymmetries of 8%
11 measured using the unilateral isometric squat. However, with limited data relating
12 specifically to asymmetries in strength and their effects on performance outcomes, a specific
13 threshold cannot be substantiated at this time. For jump testing, asymmetries > 10% have
14 been associated with a 9cm reduction in jump height (7); whereas, inter-limb differences ~10%
15 in jump height (43) and power (32) have shown minimal effect on CODS performance. This
16 provides further support for task-specificity pertaining to asymmetries, making it challenging
17 to draw definitive conclusions regarding critical thresholds during jumping-based tasks as
18 well. The relationship between asymmetry and injury has also been investigated and available
19 data indicate inter-limb differences > 15% increases the risk of injury (24,38,48). Therefore,
20 it could be suggested that a greater magnitude of asymmetry is required to place athletes at
21 heightened risk.

22 A final point to consider when comparing different asymmetry scores across different studies
23 is that multiple equations have been used to quantify inter-limb differences. Inconsistencies
24 in the equations used can result in different asymmetry percentages even when the same score
25 is attained by an athlete (10). For example, right vs. left jump height scores of 25 and 20cm

1 can result in asymmetry values between 7.04-22.2% dependent on the equation used (10). A
2 true comparison between studies would only exist if the same equation was utilized.
3 Therefore, when measuring asymmetry, practitioners are encouraged to always use the same
4 equation to ensure consistent and accurate longitudinal tracking of athletes. It is beyond the
5 scope of this review to discuss the merits of each equation; however, the interested reader is
6 encouraged to view the work of Bishop et al. (10) for further details.

7

8 **Testing Battery**

9 Based on the aforementioned evidence, a testing battery has been proposed for the assessment
10 of asymmetries in strength and jumping-based tasks (Table 2). With strength being of
11 undeniable importance in athletic performance (64), and jumping tasks occurring frequently
12 in sporting actions (29,30), testing inter-limb differences for both competencies seems logical
13 and may allow for a more complete picture of asymmetries. In addition, Table 3 provides an
14 overview of instructions for each test so that practitioners can adhere to the methods that are
15 likely to elicit the most reliable results. It should be noted that determining inter-limb
16 asymmetries during sprinting and CODS tasks would also provide S&C coaches with useful
17 information. However, the literature pertaining to asymmetries and these physical qualities is
18 scarce and further research in these areas is required before any suggestions are made.

19

20 *** *INSERT TABLES 2 AND 3 ABOUT HERE* ***

21

22 **Practical Considerations for Testing**

1 Regardless of whether asymmetries are being calculated for strength or jumping tests, there
2 are additional test considerations that practitioners should be aware of. Firstly, experience of
3 the tester must be considered. It is common for certain tests to have specific requirements that
4 aid in the standardisation of procedures. For example, it is often suggested that athletes
5 should pull “as hard and as fast as possible” when performing the IMTP test (19); therefore,
6 some level of experience or familiarity is required to know that this will likely elicit
7 favourable results in variables such as RFD especially. Secondly, the ease of testing
8 equipment must also be deliberated and it is likely that different considerations exist for
9 strength and jump tests. For example, without twin force plates it is impossible to gauge
10 information pertaining to vGRF asymmetries during strength testing. Whilst an alternative
11 solution is to test for asymmetries using isokinetic dynamometry, this method is confined to a
12 laboratory and may not be viable for many practitioners. Therefore, calculating asymmetries
13 in strength will likely require force plates. For jump tests, many alternative options exist;
14 however, force plates should still be considered a favourable option with multiple metrics
15 available for both propulsive and landing forces. Alternatively, equipment such as Optojump
16 and electronic jump mats will allow for asymmetries in metrics such as stiffness, ground
17 contact time, and jump height to be determined. Therefore, if practitioners are unable to
18 access force plates, viable alternatives do exist for jump testing in the field. Practitioners
19 constrained by budgetary restrictions require simpler and more cost-effective methods
20 whereby jump mats may be the default option. More recently, mobile technology in the form
21 of the My Jump app has also been shown to be reliable for bilateral and unilateral jump
22 testing (11). Therefore, whilst the gold standard is always preferable, measurement of
23 asymmetries during jump tests should be considered by all practitioners regardless of budgets
24 due to the wide range of options available.

1 **Conclusion**

2 The aforementioned evidence would indicate that there are advantages to choosing isometric
3 squats or the IMTP (both bilateral and unilateral variations) when quantifying asymmetries in
4 strength. Measuring peak force in particular would appear to be reliable across multiple
5 populations; and the isometric squat has shown that higher asymmetries are associated with
6 negative impacts on sport-specific tasks, and thus, performance. When combined with the
7 fact that force plates are more easily accessible in the field due to the creation of more cost-
8 effective versions, and dynamometry measures are primarily restricted to a laboratory setting,
9 the IMTP or isometric squat are the favourable options when quantifying asymmetries in
10 strength. When calculating asymmetries via jump testing, the concept of movement
11 variability must not be forgotten. Initial jump testing may be best served both bilaterally and
12 unilaterally for asymmetry detection. Once practitioners have determined the most reliable
13 and appropriate test from the battery of jump tests, this will help to streamline future test
14 protocols when determining inter-limb differences. Practitioners should keep in mind that
15 asymmetries have been frequently shown to be task-dependent and although specific
16 thresholds for reduced performance may not exist at present, asymmetries > 15% may
17 increase an athlete's risk of injury. A final thought to this review is that by assessing
18 asymmetries in both strength and jumping tasks, a more complete asymmetry picture is
19 provided by quantifying differences for two commonly prescribed exercise types in S&C
20 programming. This may allow practitioners to prioritize specific areas of their athletes'
21 training programs and target reductions in asymmetry that are relevant to the athlete.

22

23

24

1 **References**

- 2 1. Aagaard P. Training-induced changes in neural function. *Exer Sp Sci Rev* 32: 61-67,
3 2003.
- 4 2. Arteaga R, Dorado C, Chavarren J and Calbet JAL. Reliability of jumping
5 performance in active men and women under different stretch loading conditions. *J Sp*
6 *Med Phys Fit* 40: 26-34, 2000.
- 7 3. Ayala F, De Ste Croix M, Sainz de Baranda P and Santonja F. Absolute reliability of
8 isokinetic knee flexion and extension measurements adopting a prone position. *Clin*
9 *Phys Func Imaging* 33: 45-54, 2013.
- 10 4. Bailey C, Sato K, Alexander R, Chiang C-Y and Stone MH. Isometric force
11 production symmetry and jumping performance in collegiate athletes. *J Train* 2: 1-5,
12 2013.
- 13 5. Bailey C, Sato K, Burnett A and Stone MH. Force-production asymmetry in male and
14 female athletes of differing strength levels. *Int J Sports Phys Perf* 10: 504-508, 2015.
- 15 6. Bazyler CD, Bailey CA, Chiang C-Y, Sato K and Stone MH. The effects of strength
16 training on isometric force production symmetry in recreationally trained males. *J*
17 *Train* 3: 6-10, 2014.
- 18 7. Bell, DR, Sanfilippo, JL, Binkley, N and Heiderscheit, BC. Lean mass asymmetry
19 influences force and power asymmetry during jumping in collegiate athletes. *J*
20 *Strength Cond Res* 28: 884-891, 2014.
- 21 8. Benjanuvatra N, Lay BS, Alderson JA and Blanksby BA. Comparison of ground
22 reaction force asymmetry in one- and two-legged countermovement jumps. *J Str*
23 *Cond Res* 27: 2700-2707, 2013.

- 1 9. Bennell K, Wajswelner H, Lew P, Schall-Riauour A, Leslie S, Plant D and Cirone J.
2 Isokinetic strength testing does not predict hamstring injury in Australian Rules
3 footballers. *Br J Sports Med* 32: 309-314, 1998.
- 4 10. Bishop C, Read P, Chavda S and Turner A. Asymmetries of the lower limb: The
5 calculation conundrum in strength training and conditioning. *Str Cond J* 38: 27-32,
6 2016.
- 7 11. Bishop C, Tarrant J, Jarvis P and Turner A. Using the split squat to potentiate bilateral
8 and unilateral jump performance. *J Str Cond Res* Published ahead of print.
- 9 12. Bolgla LA and Keskula DR. Reliability of lower extremity functional performance
10 tests. *J Orthop Sp Phys Ther* 26: 138-142, 1997.
- 11 13. Ceroni D, Martin XE, Delhumeau C and Farpour-Lambert NJ. Bilateral and gender
12 differences during single-legged vertical jump performance in healthy teenagers. *J Str*
13 *Cond Res* 26: 452-457, 2012.
- 14 14. Cormack SJ, Newton RU, McGuigan MR and Doyle TLA. Reliability of measures
15 obtained during single and repeated countermovement jumps. *Int J Sp Phys Perf* 3:
16 131-144, 2008.
- 17 15. Costa Silva JRL, Detanico D, Dal Pupo J and Freitas C. Bilateral asymmetry of knee
18 and ankle isokinetic torque in soccer players u20 category. *Braz J Kinanthro Human*
19 *Perf* 17: 195-204, 2015.
- 20 16. Croisier J-L, Forthomme B, Namurois M-H, Vanderthommen M and Crielaard J-M.
21 Hamstring muscle strain recurrence and strength performance disorders. *Am J Sports*
22 *Med* 30: 199-203, 2002.
- 23 17. Cronin JB, Hing RD and McNair PJ. Reliability and validity of a linear position
24 transducer for measuring jump performance. *J Str Cond Res* 18: 590-593, 2004.

- 1 18. Dickin CD and Too D. Effects of movement velocity and maximal concentric and
2 eccentric actions on the bilateral deficit. *Res Quart Ex Sp* 77: 296-303, 2006.
- 3 19. Dos'Santos T, Thomas C, Jones PA and Comfort P. Assessing muscle strength
4 asymmetry via a unilateral stance isometric mid-thigh pull. *Int J Sp Phys Perf*
5 Published ahead of print.
- 6 20. Exell TA, Irwin G, Gittoes MJR and Kerwin DG. Implications of intra-limb
7 variability on asymmetry analyses. *J Sp Sci* 30: 403-409, 2012.
- 8 21. Flanagan EP, Ebben WP and Jensen RL. Reliability of the reactive strength index and
9 time to stabilisation during depth jumps. *J Str Cond Res* 22: 1677-1682, 2008.
- 10 22. Flanagan SP and Salem GJ. Bilateral differences in the net joint torques during the
11 squat exercise. *J Str Cond Res* 21: 1220-1226, 2007.
- 12 23. Fort-Vanmeerhaeghe A, Gual G, Romero-Rodriguez D and Unnitha V. Lower limb
13 neuromuscular asymmetry in volleyball and basketball players. *J Human Kin* 50: 135-
14 143, 2016.
- 15 24. Greenberger, HB and Paterno, MV. Relationship of knee extensor strength and
16 hopping test performance in the assessment of lower extremity function. *J Orthop*
17 *Sports Phys Ther* 22: 202-206, 1995.
- 18 25. Hart NH, Nimphius S, Cochrane JL and Newton RU. Reliability and validity of
19 unilateral and bilateral isometric strength measures using a customised, portable
20 apparatus. *J Aust Str Cond* 20: 61-67, 2012.
- 21 26. Hart NH, Nimphius S, Spiteri T and Newton RU. Leg strength and lean mass
22 symmetry influences kicking performance in Australian Football. *J Sp Sci Med* 13:
23 157-165, 2014.

- 1 27. Hart NH, Nimphius S, Weber J, Spiteri T, Rantalainen T, Dobbin M and Newton RU.
2 Musculoskeletal asymmetry in football athletes: A product of limb function over time.
3 *Med Sci Sp Ex* 48: 1379-1387, 2016.
- 4 28. Henry GJ, Dawson B, Lay BS and Young WB. Relationships between reactive agility
5 movement time and unilateral vertical, horizontal, and lateral jumps. *J Str Cond Res*
6 30: 2514-2521, 2013.
- 7 29. Hewit J, Cronin J and Hume P. Multidirectional leg asymmetry assessment in sport.
8 *Str Cond J* 34: 82-86, 2012.
- 9 30. Hewit JK, Cronin JB and Hume PA. Asymmetry in multidirectional jumping tasks.
10 *Phys Ther Sp* 13: 238-242, 2012.
- 11 31. Hodges SJ, Patrick RJ and Reiser RF. Effects of fatigue on bilateral ground reaction
12 force asymmetries during the squat exercise. *J Str Cond Res* 25: 3107-3117, 2011.
- 13 32. Hoffman, JR, Ratamess, NA, Klatt, M, Faigenbaum, AD and Kang, J. Do bilateral
14 power deficits influence direction-specific movement patterns? *Res Sports Med* 15: 1-
15 8, 2007.
- 16 33. Hopkins W. How to interpret changes in an athletic performance test. *Sportscience* 8:
17 1-7, 2004.
- 18 34. Impellizzeri FM, Bizzini M, Rampinini E, Cereda F and Maffiuletti NA. Reliability of
19 isokinetic strength imbalance ratios measured using the Cybex Norm dynamometer.
20 *Clin Phys Func Imaging* 28: 113-119, 2008.
- 21 35. Impellizzeri FM, Rampinini E, Maffiuletti N and Marcora SM. A vertical jump force
22 test for assessing bilateral strength asymmetry in athletes. *Med Sci Sp Ex* 39: 2044-
23 2050, 2007.
- 24 36. Jones PA and Bampouras TM. A comparison of isokinetic and functional methods of
25 assessing bilateral strength imbalance. *J Str Cond Res* 24: 1553-1558, 2010.

- 1 37. Jonhagen S, Nemeth G and Eriksson E. Hamstring injuries in sprinters: The role of
2 concentric and eccentric hamstring muscle strength and flexibility. *Am J Sports Med*
3 22: 262-266, 1994.
- 4 38. Jordan MJ, Aagaard P and Herzog W. Lower limb asymmetry in mechanical muscle
5 function: A comparison between ski racers with and without ACL reconstruction.
6 *Scandinavian J Med Sci Sports* doi: 10.1111/sms.12314
- 7 39. Joseph M, Tiberio D, Baird JL, Trojian TH, Anderson JM, Kraemer WJ and Maresh
8 CM. Knee valgus during drop jumps in National Collegiate Athletic Association
9 Division I female athletes: The effect of a medial post. *Am J Sp Med* 36: 285-289,
10 2008.
- 11 40. Kawamori N, Rossi SJ, Justice BD, Haff EE, Pistilli EE, O'Bryant HS, Stone MH and
12 Haff GG. Peak force and rate of force development during isometric and dynamic
13 mid-thigh clean pulls performed at various intensities. *J Str Cond Res* 20: 483-491,
14 2006.
- 15 41. Kobayashi Y, Kubo J, Matsubayashi T, Matsuo A, Kobayashi K and Ishii N.
16 Relationship between bilateral differences in single-leg jumps and asymmetry in
17 isokinetic knee strength. *J App Biomech* 29: 61-67, 2013.
- 18 42. Kristianslund E and Krosshaug T. Comparison of drop jumps and sport-specific
19 sidestep cutting: Implications for anterior cruciate ligament injury risk screening. *Am*
20 *J Sp Med* 41: 684-688, 2013.
- 21 43. Lockie RG, Callaghan SJ, Berry SP, Cooke ERA, Jordan CA, Luczo TM and Jeffriess
22 MD. Relationship between unilateral jumping ability and asymmetry on
23 multidirectional speed in team-sport athletes. *J Str Cond Res* 28: 3557-3566, 2014.
- 24 44. Maloney SJ, Fletcher IM and Richards J. A comparison of methods to determine
25 bilateral asymmetries in vertical leg stiffness. *J Sports Sci* 34: 829-835, 2016.

- 1 45. Maloney SJ, Richards J, Nixon DGD, Harvey LJ and Fletcher IM. Do stiffness and
2 asymmetries predict change of direction performance? *J Sports Sci* DOI:
3 <http://dx.doi.org/10.1080/02640414.2016.1179775>
- 4 46. Meylan C, McMaster T, Cronin J, Mohammad NI, Rogers C and deKlerk M. Single-
5 leg lateral, horizontal and vertical jump assessment: Reliability, interrelationships, and
6 ability to predict sprint and change-of-direction performance. *J Str Cond Res* 23:
7 1140-1147, 2009.
- 8 47. Newton, RU, Gerber, A, Nimphius, S, Shim, JK, Doan, BK, Robertson, M, Pearson,
9 DR, Craig, BW, Hakkinen, K and Kraemer, WJ. Determination of functional strength
10 imbalance of the lower extremities. *J Strength Cond Res* 20: 971-977, 2006.
- 11 48. Noyes FR, Barber SD and Mangine RE. Abnormal lower limb symmetry determined
12 by function hop tests after anterior cruciate ligament rupture. *Am J Sp Med* 19: 513-
13 518, 1991.
- 14 49. Pain MTG. Considerations for single and double leg drop jumps: Bilateral deficit,
15 standardizing drop height, and equalising training load. *J App Biomech* 30: 722-727,
16 2014.
- 17 50. Pappas E and Carpes FP. Lower extremity kinematic asymmetry in male and female
18 athletes performing jump-landing tasks. *J Sci Med Sport* 15: 87-92, 2012.
- 19 51. Rannama I, Port K, Bazanov B and Pedak K. Sprint cycling performance and
20 asymmetry. *J Human Sport Ex* 10: 247-258, 2015.
- 21 52. Reid A, Birmingham TB, Stratford PW, Alcock GK and Giffin JR. Hop testing
22 provides a reliable and valid outcome measure during rehabilitation after anterior
23 cruciate ligament reconstruction. *Phys Ther* 87: 337-349, 2007.


- 1 53. Ritti-Dias RM, Avelar A, Salvador EP and Cyrino ES. Influence of previous
2 experience on resistance training on reliability of one-repetition maximum test. *J Str*
3 *Cond Res* 25: 1418-1422, 2011.
- 4 54. Rohman E, Steubs JT and Tompkins M. Changes in involved and uninvolved limb
5 function during rehabilitation after anterior cruciate ligament reconstruction:
6 Implications for limb symmetry index measures. *Am J Sp Med* 43: 1391-1398, 2015.
- 7 55. Ross MD, Langford B and Whelan PJ. Test-retest reliability of 4 single-leg horizontal
8 hop tests. *J Str Cond Res* 16: 617-622, 2002.
- 9 56. Ruas CV, Brown LE and Pinto RS. Lower-extremity side-to-side strength asymmetry
10 of professional soccer players according to playing position. *Kinesiology* 2: 188-192,
11 2015.
- 12 57. Salonikidis K, Amiridis IG, Oxyzoglou N, de Villareal ES, Zafeiridis A and Kellis E.
13 Force variability during isometric wrist flexion in highly skilled and sedentary
14 individuals. *Euro J App Phys* 107: 715-722, 2009.
- 15 58. Sato, K and Heise, GD. Influence of weight distribution asymmetry on the
16 biomechanics of a barbell squat. *J Strength Cond Res* 26: 342-349, 2012.
- 17 59. Schiltz M, Lehance C, Maquet D, Bury T, Crielaard J-M and Croisier J-L. Explosive
18 strength imbalances in professional basketball players. *J Ath Train* 44: 39-47, 2009.
- 19 60. Seo D, Kim E, Fahs CA, Rossow L, Young K, Ferguson SL, Thiebaud R, Sherk VD,
20 Loenneke JP, Kim D, Lee M, Choi K, Bemben DA, Bemben MG and So W-Y.
21 Reliability of the one-repetition maximum test based on muscle group and gender. *J*
22 *Sp Sci Med* 11: 221-225, 2012.
- 23 61. Sheppard JM and Young WB. Agility literature review: Classifications, training and
24 testing. *J Sports Sci* 24: 919-932, 2006.

- 1 62. Stalboom M, Holm DJ, Cronin J and Keogh J. Reliability of kinematics and kinetics
2 associated with single leg drop jump assessment. A brief report. *J Sp Sci Med* 6: 261-
3 264, 2007.
- 4 63. Stephens II TM, Lawson BR, DeVoe DE and Reiser II RF. Gender and bilateral
5 differences in single-leg countermovement jump performance with comparison to a
6 double-leg jump. *J App Biomech* 23: 190-202, 2007.
- 7 64. Suchomel TJ, Nimphius S and Stone MH. The importance of muscular strength in
8 athletic performance. *Sports Med* [Epub. ahead of print]:1-31, 2016.
- 9 65. Sugiyama T, Kameda M, Kageyama M, Kiba K, Kanehisa H and Maeda A.
10 Asymmetry between the dominant and non-dominant legs in the kinematics of the
11 lower extremities during a running single leg jump in collegiate basketball players. *J*
12 *Sports Sci Med* 13: 951-957, 2014.
- 13 66. Turner A, Brazier J, Bishop C, Chavda S, Cree J and Read P. Data analysis for
14 strength and conditioning coaches: Using excel to analyse reliability, differences, and
15 relationships. *Str Cond J* 37: 76-83, 2015.
- 16 67. Walsh MS, Ford KR, Bangen KJ, Myer GD and Hewett TE. The validation of a
17 portable force plate for measuring force-time data during jumping and landing tasks. *J*
18 *Str Cond Res* 20: 730-734, 2006.
- 19 68. Weir JP. Quantifying test-retest reliability using the intraclass correlation coefficient
20 and the SEM. *J Str Cond Res* 19: 231-240, 2005.
- 21 69. Yoshioka S, Nagano A, Hay DC and Fukashiro S. The effect of bilateral asymmetry
22 of muscle strength on jumping height of the countermovement jump: A computer
23 simulation study. *J Sp Sci* 28: 209-218, 2010.
- 24 70. Zifchock R, Davis I, Higginson J and Royer T. The symmetry angle: A novel, robust
25 method of quantifying asymmetry. *Gait & Posture* 27: 622-627, 2008.

1 Table 1: Intraclass correlation coefficients (ICC) and coefficient of variations (CV) for peak
 2 force, mean force, and RFD during bilateral and unilateral isometric squats (adapted from
 3 Hart et al. [25])

Variable		Bilateral ISO Squat	Unilateral ISO Squat (D)	Unilateral ISO Squat (ND)
Peak Force	ICC	0.97	0.96	0.98
	CV (%)	3.6	4.7	3.6
Mean Force	ICC	0.91	0.95	0.83
	CV (%)	8.4	6.1	9.3
RFD	ICC	0.94	0.93	0.36
	CV (%)	15.2	14.5	45.5
D = Dominant limb, ND = Non-dominant limb, RFD = Rate of force development				

4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19



9 Figure 1: Diagrammatic representation of four commonly used hop tests to determine
 10 asymmetries. The single leg hop requires one maximal jump landing on the same limb.
 11 Failure to land without falling over or ‘bouncing forward’ requires the test to be retaken. The
 12 triple hop assesses maximal distance for three hops in a rebounding pattern. A stable landing
 13 must also be demonstrated for the final hop. The 6m timed hop positions timing gates at 0
 14 and 6m and asks subjects to hop on one limb as fast as they can for the total distance; thus,
 15 reporting an outcome of time. The crossover hop requires three maximal hops (for distance)
 16 in a diagonal pattern. A stable landing must also be demonstrated on the final hop.

17
 18
 19
 20
 21
 22

1 Table 2: Proposed testing battery for the assessment of asymmetries

	Metrics	Selected Test	Testing Equipment
Strength tests	Peak force, mean force, RFD, impulse	IMTP or isometric squat (+ SL variations)	Force plates
Jump tests	Peak force, impulse, jump height or distance	CMJ and BJ (+ SL variations)	Force plates (or jump mat), measuring tape
RFD = Rate of force development, IMTP = Isometric mid-thigh pull, SL = Single leg, CMJ = Countermovement jump, BJ = Broad jump			

- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15

1 Table 3: Instructions for how to administer different asymmetry tests

Test	Procedural Instructions
IMTP	<p>Previous literature has outlined the knee angle to be set at 125° and the hip angle at 175° (4), with 180° representing full extension at both joints. Joint angles can be measured manually using a goniometer and weightlifting straps can be used to ensure a more secure grip on the bar. Once the position is assumed, athletes should be instructed to pull “as hard and as fast as possible” (16) which may aid in producing reliable results for variables such as RFD when measuring on force plates. For the unilateral version of this test, Dos Santos et al. (16) suggested that the non-stance limb be flexed 90° at the knee joint.</p>
Isometric Squat	<p>Hip and knee angles should be set at 140° with the bar resting on the upper trapezius muscle (as per standard high-bar back squat technique) (23,24). Athletes should be instructed to push “as hard and as fast as possible” which may aid in producing reliable results for variables such as RFD when measuring on force plates. For the unilateral version of this test, although not specified by Hart et al. (23,24), it seems logical to ask athletes to flex their non-working limb’s knee joint to 90°, as suggested for the IMTP procedures.</p>
CMJ	<p>Hands should be fixed onto hips so as to minimise any contribution from the upper body. Upon instruction, the athlete can dip to a self-selected depth during the countermovement prior to accelerating vertically as explosively as possible. Lower limbs should remain extended at all times during the flight phase of the jump before landing back on the force plate, jump mat, or ground. The same procedures should be followed for unilateral versions of this test.</p>
BJ	<p>Hands should be fixed onto hips so as to minimise any contribution from the upper body. Upon instruction, the athlete can dip to a self-selected depth during the countermovement prior to accelerating horizontally as explosively as possible, with the aim being to jump as far as possible (i.e.: a standing long jump). Trials are void and must be repeated if athletes are unable to stabilise on landing. When measuring distance, the reading should be taken (to the nearest millimetre) from the rear most heel closest to the start position. The same procedures should be followed for unilateral versions of this test.</p>

IMTP = Isometric mid-thigh pull, RFD = Rate of force development,
CMJ = Countermovement jump, BJ = Broad jump

1

2