

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Aceijas, Carmen ORCID logoORCID: <https://orcid.org/0000-0002-3652-6536>, Bello-Corassa, Rafael, Waldhäusl, Sabrina, Lambert, Nicky ORCID logoORCID: <https://orcid.org/0000-0001-8785-4719> and Cassar, Simon (2016) Barriers and determinants of physical activity among UK university students. In: 9th EPH conference. All for Health - Health for All, 9 - 12 November, 2016, Vienna, Austria. . [Conference or Workshop Item]

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/22627/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

BARRIERS AND DETERMINANTS OF PHYSICAL ACTIVITY AMONG UK UNIVERSITY STUDENTS

Aceijas C^a, Bello-Corassa R^b, Waldhäusl S^c, Lambert N^d, Cassar S^e

^a Corresponding author. Natural Sciences. School of Science and Technology. Middlesex University. London. UK. Email address: c.aceijas@mdx.ac.uk

^b Universidade Federal dos Vales do Jequitinhonha e Mucuri. Alto da Jacuba. Brazil. Email address: rafael.bellocorassa@gmail.com

^c Natural Sciences. School of Science and Technology. Middlesex University. London. UK. Email address: SW1123@live.mdx.ac.uk

^d Mental Health and Social Work. School of Health and Education. Middlesex University. London. UK. Email address: s.cassar@mdx.ac.uk

^e Wellbeing Services. Middlesex University London. UK n.lambert@mdx.ac.uk

Background: Association between high physical activity (PA) levels and lower health risks and between sedentary behaviours and higher health risks have been consistently identified in scientific literature. This study investigated determinants of physical activity among UK university students.

Methods: Online Survey among Middlesex University London UG students with stratified random sampling strategy (n=468) to assess students' PA levels. Stratified by gender bivariate analysis and multivariate logistic regression were used identify risk factors of suboptimal PA. Additionally, two students' focus groups and three university stakeholders in-depth interviews, were conducted.

Results: 60% of respondents were insufficiently active. 89% do not use the university gym due to lack of time (40%), price (30%), and embarrassment (11%). PA was significantly associated with Gender – woman had lower PA levels (OR=2.8; 95%CI =1.2–6.2); Ethnicity – black and mixed ethnicity students had lower and higher PA levels, respectively; not using the university gym (OR=2.8; 95%CI=1.2–6.3). Stratified bivariate analysis showed that underweight and obese women had lower PA levels; disabled men had a fivefold risk for insufficient PA (OR=5.1; 95%CI=1.03–24.92); males living with their parents during term were more active. Multivariate model withheld three variables, gender, gym use and smoking – smokers were less active (OR=2.1; 95%CI=1.0–4.3). Qualitative results showed finances, academic pressure and university systems as barriers for PA.

Conclusions: PA is associated with various factors, gender being the most important. Regular moderate/vigorous PA appears as protective factor against smoking. Lack of time and cost were institutional barriers for PA. Universities must implement strategies to reduce cost, increase accessibility and improve students' time management capacity to include PA in their routines. Furthermore, tailor made and outside gym sessions may help in achieving optimal PA.

Main message:

Students' PA levels are low and likely to compromise their health. Universities are in a strategic position to influence their choices and should incorporate strategies to increase their PA levels.