

# Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

McCubbine, Jermaine, Turner, Anthony N., Dos'Santos, Thomas and Bishop, Chris (2018)  
Reliability and measurement of inter-limb asymmetries in four unilateral jump tests in elite youth  
female soccer players. *Professional Strength & Conditioning* (49). pp. 7-12. ISSN 1757-5834

Final accepted version (with author's formatting)

This version is available at: <http://eprints.mdx.ac.uk/23038/>

## Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

[eprints@mdx.ac.uk](mailto:eprints@mdx.ac.uk)

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

1 RELIABILITY AND MEASUREMENT OF INTER-LIMB ASYMMETRIES  
2 IN 4 UNILATERAL JUMP TESTS IN ELITE YOUTH FEMALE SOCCER  
3 PLAYERS

---

4

5

6

7 **AUTHORS:**

8 Jermaine McCubbine <sup>1</sup> (MSc, ASCC), Anthony Turner <sup>2</sup> (PhD, ASCC, CSCS\*D), Tom Dos'  
9 Santos <sup>3</sup> (MSc) and Chris Bishop <sup>2</sup> (MSc, ASCC)

10

11

12 **AFFILIATIONS:**

13 1. Arsenal Football Club Training Ground, Bell Lane, London Colney, St. Albans, AL2  
14 1DR, UK

15 2. London Sport Institute, School of Science and Technology, Middlesex University,  
16 Greenlands Lane, Allianz Park, NW4 1RL, London, UK

17 3. Directorate of Sport, Exercise and Physiotherapy, University of Salford, Frederick  
18 Road, Salford, Greater Manchester M6 6PU, UK

19

20

21 **CORRESPONDENCE:**

22 **Name:** Chris Bishop

23 **Email:** [C.Bishop@mdx.ac.uk](mailto:C.Bishop@mdx.ac.uk)

24 **Address:** As above for affiliation 1.

25 RELIABILITY AND MEASUREMENT OF INTER-LIMB ASYMMETRIES  
26 IN 4 UNILATERAL JUMP TESTS IN ELITE YOUTH FEMALE SOCCER  
27 PLAYERS

---

28 **ABSTRACT**

29 **Purpose:** The purpose of this study was to determine the within and between-session  
30 reliability, and inter-limb asymmetries, in four unilateral jump tests in elite youth female  
31 soccer players. Given the low plyometric training age and paucity of data for this population,  
32 this research study was warranted. **Methods:** Nineteen elite youth female soccer players (age:  
33  $10 \pm 1.1$  years; height:  $141 \pm 7.9$  cm; body mass:  $35 \pm 7.1$  kg) were recruited from an elite  
34 Tier 1 Regional Talent Centre of a professional soccer club. Tests included the single leg  
35 countermovement jump (SLCMJ), single leg hop, triple hop, and crossover hops for distance  
36 with reliability quantified via the coefficient of variation (CV), intraclass correlation  
37 coefficient (ICC), and standard error of the measurement (SEM). Inter-limb asymmetries  
38 were also calculated. **Results:** Both test sessions resulted in excellent within-session  
39 reliability (ICC range = 0.81-0.99; SEM range = 0.11-0.49; and CV range = 2.6-6.0%).  
40 Between-session reliability was deemed good to excellent (ICC range = 0.72-0.99 and pooled  
41 CV = 2.7-5.7%). Asymmetries were deemed small across both test sessions with the highest  
42 value reported in the SLCMJ (6.12%). **Conclusion:** Results highlight that unilateral jump  
43 tests can be considered a reliable test protocol in elite youth female soccer players, which is  
44 important considering youth athletes likely do not have a vast plyometric training age.  
45 Furthermore, inter-limb differences appear small in the present sample which may also be  
46 explained by their limited training age, given that asymmetries have previously been  
47 highlighted to be a product of limb function over time.

48 **Key Words:** Lower extremity, single leg, youth athletes

49 **INTRODUCTION**

50 Physical performance testing is a common component for strength and conditioning  
51 practitioners to undertake, allowing athletes' fitness capabilities to be effectively monitored  
52 which in turn may aid the decision-making process during the design of training  
53 interventions. Numerous factors must be considered when selecting appropriate fitness tests  
54 to measure performance including: age, equipment, environment, time, training age, and the  
55 reliability of the test itself need to be determined in order to assess if a given protocol is to be  
56 included or excluded from a test battery (5). Test reliability is crucial as this will enable  
57 practitioners to determine if the chosen assessment produces consistent results; thus, allowing  
58 results to be interpreted with confidence (33).

59 Jump testing is a common mode of assessment as it provides a relatively quick and reliable  
60 method for assessing the explosive capabilities of athletes (20,22,34). Sports such as soccer,  
61 basketball and volleyball, have used bilateral countermovement jumps (CMJ), drop jumps  
62 (DJ), and squat jumps (SJ) to assess lower limb power (13,26,35). However, many team sport  
63 actions (such as jumping, changing direction, and sprinting) occur unilaterally; thus, this  
64 provides additional considerations for test protocols taking place on one leg (5,30).  
65 Furthermore, given the multi-planar nature of team sports (12), testing protocols should  
66 reflect this also so that they can be considered ecologically valid for the population in  
67 question. Consequently, a variety of unilateral jump tests exist that enable the aforementioned  
68 factors to be accounted for. However, owing to the high degree of movement variability  
69 associated with jump testing (17), and the heightened instability of testing unilaterally,  
70 assessing the reliability of these tests becomes even more important. The single leg  
71 countermovement jump (SLCMJ) and a variety of hop tests (single leg hop, triple hop, and  
72 crossover hop) have been commonly used in the literature. Typically, test-retest reliability of  
73 these tests appears strong with intraclass correlation coefficient (ICC) values between 0.8-

74 0.98 (7,14,21,23,25,28,29,31). However, considerably less information exists about the  
75 reliability of these tests for youth athletes (9,24), especially those competing at the highest  
76 level in female youth soccer. Given that youth athletes likely have a lower plyometric  
77 training age than adults, additional reliability data on this population is warranted.

78 An additional advantage of selecting unilateral variations of these jump tests is that they also  
79 enable inter-limb asymmetries to be quantified. Recent literature has highlighted that  
80 unilateral jump testing is a useful method for quantifying between-limb differences (3,4,5). In  
81 addition, it has been suggested that assessing asymmetries from unilateral tests may be more  
82 applicable than their bilateral counterparts because no contribution from the other limb is  
83 present (2,5). This is compounded from research by Jordan et al. (17), who highlighted the  
84 changing nature of asymmetries during different phases of bilateral jumping before take-off.  
85 From a physical performance perspective, Maloney et al. (19) reported that inter-limb  
86 differences in jump height were an important factor in explaining slower change of direction  
87 times and additional negative associations between strength asymmetries and performance  
88 have also been noted (4). When considering injury risk, asymmetries of 15% have historically  
89 been suggested as a threshold to be mindful of (15,28). However, more recent research has  
90 proposed that patients are four times more likely to re-rupture their anterior cruciate ligament  
91 (ACL) if an asymmetry threshold of 10% is not met from hop testing as part of a return to  
92 sport criteria (18). Thus, the presence of inter-limb differences may have implications on  
93 physical performance and injury risk, indicating their quantification from unilateral jump  
94 testing is warranted.

95 Therefore, the aims of the present study were to report within and between-session reliability  
96 of four commonly used unilateral jump tests (SLCMJ, single leg hop, triple hop, and  
97 crossover hop for distance) in elite youth female soccer players. In addition, inter-limb  
98 differences were also quantified for all jump tests enabling the creation of a multi-directional

99 asymmetry profile for this population.

100

## 101 **METHODS**

### 102 **Subjects**

103 Nineteen elite youth female soccer players (age:  $10 \pm 1.1$  years; height:  $141 \pm 7.9$  cm; body  
104 mass:  $35 \pm 7.1$  kg), were recruited from a Tier 1 Regional Talent Centre (RTC) of a  
105 professional soccer club. Subjects were considered elite as this particular level is the highest  
106 standard of female youth club soccer in England. Players trained for at least 36 weeks per  
107 year and were required to partake in a minimum of one hour of structured strength and  
108 conditioning training per week. Emphasis at this age was placed on mastering fundamental  
109 movement patterns, building strong foundations, enhancing technical competency, and  
110 improving general motor control. All subjects were free from injury and any player who  
111 presented an injury resulting in more than one week of missed training prior to testing was  
112 excluded. Informed consent and PAR-Q forms were completed from all relevant  
113 parent/guardians as all participants were under the age of 18. Ethical approval was granted  
114 from the London Sports Institute ethics committee, Middlesex University.

### 115 **Procedures**

116 Participants were tested at the same time of day on three separate testing occasions, each  
117 separated by seven days. Session one was used to familiarize all participants with the test  
118 procedures, allowing them to practice each jump test as many times as they wanted. A  
119 particular emphasis was placed on landing mechanics, owing to the increased demand of  
120 having to land on one limb in the chosen tests. The next two sessions were used for official  
121 testing and data collection. All participants were asked not to participate in any strenuous

122 exercise at least 24 hours prior to testing, and to ensure they wore the same footwear on each  
123 occasion to negate the effects of different shoe design and support structures. Both testing  
124 sessions took place on a third generation pitch, which subjects were used to training on twice  
125 weekly. Each test consisted of three trials on each limb with 60 seconds rest between trials,  
126 and 2-minutes rest between tests, to enable full recovery (28). All tests were conducted in a  
127 randomized, counter-balanced order, to negate any potential learning effects. Before  
128 familiarization and testing sessions, all participants completed a standardized warm-up  
129 protocol (Table 1), following the RAMP system as outlined by Jeffreys (16). This consisted  
130 of dynamic exercises progressing from low intensity and generic movements to higher  
131 intensity with more specific movement patterns. A 3-minute rest period was prescribed  
132 between the completion of the warm up and commencement of the first test.

133

134 \*\*\* INSERT TABLE 1 ABOUT HERE \*\*\*

135

136 *Single leg countermovement jump (SLCMJ)*. Subjects stood in an upright position, hands on  
137 hips, with feet positioned hip width apart. One leg lifted off the floor to approximately mid  
138 shin height of the standing leg. Subjects then squatted to a self-selected depth followed by a  
139 quick upward vertical movement, jumping as high as possible. The jumping leg had to remain  
140 fully extended and hands fixed to hips; any deviation from this resulted in a retrieval after a 60-  
141 second rest period. Jump height was calculated by the flight time method using the “My  
142 Jump” iPhone application, which has been shown to be a reliable method for quantifying this  
143 outcome measure (1).

144

145 *Single Hop (for distance)*. Subjects begin by standing on a designated testing leg with hands

146 on hips and their toes behind the starting line. Subjects were then instructed to hop as far  
147 forward as possible and land on the same leg (Figure 1). Upon landing, participants were  
148 required to ‘hold and stick’ their position for two seconds. Failure to stick the landing  
149 resulted in a void trial and a retrial after a 60-second rest. This was consistent across all trials  
150 for all hop tests. The distance hopped from the starting line to the point where the subject’s  
151 landing heel hit in the final position was then recorded to the nearest centimeter using a  
152 standard measuring tape (also used for all hop tests).

153 *Triple Hop (for distance)*. Subjects begin by standing on the designated testing leg, hands on  
154 hips with their toes behind the starting line. Subjects were instructed to take three maximal  
155 hops forward, landing on the same leg throughout and holding and sticking the 3<sup>rd</sup> contact for  
156 two seconds (Figure 1). The distance hopped from the starting line to the landing position of  
157 the subjects’ heel of the same limb was then measured and recorded to the nearest centimeter.

158

159 *Crossover Hop (for distance)*. Subjects began by standing on the designated testing leg, with  
160 their toes behind the starting line. If subjects were hopping with their right leg, they started  
161 the test on the right side of the measuring tape and vice versa if they started on the left limb.  
162 Subjects were instructed to take three consecutive maximal hops forward; each time crossing  
163 over an area measuring 15 cm wide landing on the same leg throughout (Figure 1). As per  
164 previous hop testing protocols, all subjects were required to stick the final landing for two  
165 seconds. The distance hopped from the starting line to the point where the subject’s heel hit  
166 on completion of the third jump was measured and recorded to the nearest centimeter.

167

168 \*\*\* INSERT FIGURE 1 ABOUT HERE \*\*\*

169

170

171 **Statistical Analyses**

172 All data was initially computed as means and standard deviations (SD) in Microsoft Excel™.  
173 The coefficient of variation (CV) and standard error of the measurement (SEM) were used to  
174 quantify absolute reliability, whilst the intraclass correlation coefficient (ICC) with absolute  
175 agreement quantified relative reliability. Interpretation of ICC values was in accordance with  
176 previous research where values > 0.75 are considered ‘excellent’, 0.4-0.75 are considered  
177 ‘good’, and anything < 0.4 is considered ‘poor’ (10), and CV’s were deemed acceptable if <  
178 10% (6). Both within and between-session reliability were calculated with the CV being  
179 quantified in Microsoft Excel™ and all other statistics computed in SPSS (SPSS Inc.,  
180 Chicago, IL, USA). In addition, the smallest worthwhile change (SWC) was calculated by  
181 multiplying the pooled SD by 0.2 (33), and then converted to a percentage. Finally, inter-limb  
182 asymmetries were calculated using the equation: (maximum value – minimum  
183 value)/maximum value x 100 which has been previously used in research for youth soccer  
184 athletes (27), and was quantified from an average of the three trials.

185

186 **RESULTS**

187 Within-session reliability for the first testing session was excellent (Table 2: ICC range =  
188 0.82-0.99; SEM range = 0.11-0.42; and CV range = 2.6-6.0%). Within-session reliability for  
189 testing session two was also excellent (Table 3: ICC range = 0.81-0.99; SEM range = 0.16-  
190 0.49; and CV range = 2.8-5.4%). Between-session reliability for all hop tests was good to  
191 excellent (Table 4: ICC range = 0.72-0.99; pooled CV range = 2.7-5.7%; and SWC range =  
192 2.35-5.95%). Inter-limb differences were ≤ 6.12% during both sessions for all tests (Tables 2  
193 and 3).

194

195 \*\*\* INSERT TABLES 2-4 ABOUT HERE \*\*\*

196

## 197 **DISCUSSION**

198 The aims of the present study were to determine within and between-session reliability, and  
199 quantify inter-limb asymmetries from four unilateral jump tests in elite youth female  
200 soccer players. Given the paucity of data about both reliability and asymmetries in this  
201 population, this research study was warranted.

202 Tables 2 and 3 highlight within-session reliability data for both data collection sessions.  
203 Cormack et al. (6) suggest that acceptable typical error (CV) values should fall below 10%  
204 and with the highest CV value of 6.0% reported across both sessions; within-session  
205 reliability was good to excellent. In general, the SLCMJ provided the smallest typical error  
206 (2.6-3.5%) and the crossover hop the largest (3.6-6.0%). Although anecdotal, when the  
207 repeated nature of the crossover hop is combined with the inclusion of some lateral  
208 movement, it is unsurprising that larger variability is seen within this test. However, all  
209 values are still deemed to be acceptable. A similar trend is followed when interpreting results  
210 from the ICC with the reliability of the SLCMJ near perfect (0.99). In contrast, the crossover  
211 hop showed ICC values of 0.82-0.83 on the right limb; however, reliability was notably better  
212 on the left side (0.93-0.94). Regardless, with results being interpreted in line with suggestions  
213 by Fleiss, (10), all values are still considered excellent.

214 Table 4 portrays between-session reliability results, highlighting that all jump tests had good  
215 to excellent levels of reliability when using the ICC (0.72-0.96). The single leg hop was  
216 shown to be one of the least reliable test between testing sessions (ICC = 0.72-0.76), perhaps  
217 indicating that there may have been a slightly larger learning effect compared to the other  
218 protocols. However, it should be acknowledged that the only value considered 'good' was  
219 close to the 0.75 threshold needed to be classed as 'excellent'. Pooled CV values followed a  
220 similar trend to the within-session results with the SLCMJ showing the greatest consistency

221 (2.7-3.2%) and the crossover hop showing the greatest variability (4.0-5.7%). However, all  
222 results should be interpreted with confidence which is important for this age group given their  
223 lack of experience with plyometric training. Furthermore, any recorded data from such tests  
224 can be used to monitor progress over time (33) or as outcome measures to assess the  
225 effectiveness of targeted training interventions (3).

226 Tables 2 and 3 also show inter-limb asymmetry data with results highlighting small between-  
227 limb differences for all tests ( $\leq 6.12\%$ ). Despite these small values, the SLCMJ would appear  
228 to demonstrate notably larger asymmetries (5.10-6.12%) than any of the horizontal hop tests  
229 (0.24-2.02%). When interpreting asymmetry data, it is essential to understand that an inter-  
230 limb difference can only be classified as 'real' if the value is greater than the intra-limb  
231 variability (8), which in the present study is represented by the typical error (CV). The only  
232 test to exhibit real asymmetries was the SLCMJ whereas between-limb differences in all  
233 other tests can be considered as natural variability during the testing process (3).  
234 Asymmetries have been suggested as being a by-product of repeated sporting actions that  
235 occur over time (11), and with a young population tested in the present study, it is possible  
236 that their training age was too low to impact any limb dominance issues, manifested during  
237 horizontal jump testing. Furthermore, although the SLCMJ highlighted real side-to-side  
238 differences, the values can still be considered small with asymmetry and injury literature  
239 suggesting values between 10-15% as being thresholds to be aware of (15,18,28). From a  
240 longitudinal perspective, practitioners should continue to monitor inter-limb asymmetries to  
241 ensure that these values and the natural variability during testing remain small (3,5). In  
242 addition, a sport such as soccer is characterised by limb dominance (12), so increased  
243 exposure to playing and training time may be a potential cause to increase inter-limb  
244 differences if they are not monitored closely and programmes manipulated accordingly.  
245 Given recent literature highlighted that larger asymmetries may be detrimental to

246 performance (4), it is likely that monitoring asymmetries from a young age is a worthwhile  
247 process for practitioners.

248

## 249 **PRACTICAL APPLICATIONS**

250 The results from the present study highlight that unilateral jump tests are a reliable testing  
251 protocol for elite youth female soccer athletes, which is useful information given the lack of  
252 data in this population and their associated reduced plyometric training age. Inter-limb  
253 asymmetries appear small in this population which may be a by-product of a lower training  
254 age compared to adults. Practitioners can use this information to confidently incorporate  
255 unilateral jump testing with youth female athletes. However, it is still suggested that  
256 practitioners employ similar methods to quantify reliability data for their own athletes given  
257 test consistency may vary between populations. A final note of consideration is that if only  
258 one test was selected (due to time-constraints for example); the SLCMJ may be considered as  
259 the preferred choice for practitioners due to its ability to expose greater inter-limb  
260 asymmetries and stronger reliability than the horizontal hop tests.

261

262

263

264

265

266

267

268

269

270

271 **REFERENCES**

- 272 1. Balsalobre-Fernandez C, Tejero-Gonzalez CM, Del Campo- Vecino J, and Bavaresco  
273 N. The concurrent validity and reliability of a low-cost, high-speed camera-based  
274 method for measuring the flight time of vertical jumps. **J Strength Cond Res** 28:  
275 528–533, 2014.
- 276 2. Benjanuvatra N, Lay BS, Alderson JA and Blanksby BA. Comparison of ground  
277 reaction force asymmetry in one- and two-legged countermovement jumps. **J**  
278 **Strength Cond Res** 27: 2700-2707, 2013.
- 279 3. Bishop C, Read P, Chavda S, and Turner, A. 2016. Asymmetries of the lower limb:  
280 the calculation conundrum in strength training and conditioning. **Strength Cond J** 38:  
281 27-32, 2016.
- 282 4. Bishop C, Turner A, and Read P. Effects of inter-limb asymmetries on physical and  
283 sports performance: A systematic review. **J Sports Sci** Published ahead of print.  
284 <http://dx.doi.org/10.1080/02640414.2017.1361894>.
- 285 5. Bishop C, Turner A, Jarvis P, Chavda S, and Read P. Considerations for selecting  
286 field-based strength and power fitness tests to measure asymmetries. **J Strength**  
287 **Cond Res** 31: 2635-2644, 2017.
- 288 6. Cormack S, Newton R, McGuigan M, and Doyle T. Reliability of measures obtained  
289 during single and repeated countermovement jumps. **Int J Sports Phys Perf** 3: 131-  
290 144, 2008.
- 291 7. Dos'Santos T, Thomas C, Jones P, and Comfort P. Asymmetries in single and triple  
292 hop are not detrimental to change of direction speed. **J Trainology** 6: 35-41, 2017.
- 293 8. Exell T, Gittoes M, Irwin G, and Kerwin D. 2012. Gait asymmetry: Composite scores  
294 for mechanical analyses of sprint running. **J Biomechanics** 45: 1108-1111, 2012.
- 295 9. Fernandez-Santos J, Ruiz J, Cohen D, Gonzalez-Montesinos J, and Castro-Piñero J.

- 296 2015. Reliability and validity of tests to assess lower-body muscular power in  
297 children. **J Strength Cond Res** 29: 2277-2285, 2015.
- 298 10. Fleiss J. 1999. Reliability of measurement. *The Design and Analysis of Clinical*  
299 *Experiments*. Toronto, John Wiley and Son 1986;1-32.
- 300 11. Hart N, Nimphius S, Weber J, Spiteri T, Rantalainen T, Dobbin M, and Newton R.  
301 Musculoskeletal asymmetry in football athletes: a product of limb function over  
302 time. **Med Sci Sports Exer** 48: 1379-1387, 2016.
- 303 12. Hewit J, Cronin J, and Hume P. Multidirectional leg asymmetry assessment in sport.  
304 **Strength Cond J** 34: 82–86, 2012.
- 305 13. Hoff J, and Helgerud J. Endurance and strength training for soccer players:  
306 Physiological considerations. **Sports Med** 34: 165–180, 2004.
- 307 14. Hooper M, Goh C, Wentorth A, Chan K, Chau W, Wotton J, Strauss R, and Boyle.  
308 Test re-test reliability of knee rating scales and functional hop tests one year  
309 following anterior cruciate ligament reconstruction. **Phys Ther Sport** 3: 10-18, 2002.
- 310 15. Impellizzeri FM, Rampinini E, Maffiuletti N and Marcora SM. A vertical jump force  
311 test for assessing bilateral strength asymmetry in athletes. **Med Sci Sports Ex** 39:  
312 2044-2050, 2007.
- 313 16. Jeffreys I. Warm-up revisited: The ramp method of optimizing warm-ups. **Prof**  
314 **Strength Cond J** 6: 12-18, 2007.
- 315 17. Jordan M, Aagaard P, and Herzog W. Lower limb asymmetry in mechanical muscle  
316 function: A comparison between ski racers with and without ACL reconstruction.  
317 **Scand J Med Sci Sports** doi: 10.1111/sms.12314.
- 318 18. Kryitsis P, Bahr R, Landreau P, Miladi R, and Witvrouw E. Likelihood of ACL graft  
319 rupture: Not meeting six clinical discharge criteria before return to sport is associated  
320 with a four times greater risk of rupture. **Brit J Sports Med** 50: 946-951, 2016.

- 321 19. Maloney S, Richards J, Nixon D, Harvey L, and Fletcher I. Do stiffness and  
322 asymmetries predict change of direction performance? **J Sports Sci** 35: 547-556,  
323 2017.
- 324 20. Markovic G, Dizdar D, Jukic I, and Cardinale M. Reliability and factorial validity of  
325 squat and countermovement jump tests. **J Strength Cond Res** 18: 551–555, 2004.
- 326 21. Maulder P, and Cronin J. Horizontal and vertical jump assessment: reliability,  
327 symmetry, discriminative and predictive ability. **Phys Ther Sport** 6: 74-82, 2005.
- 328 22. McMaster D, Gill N, Cronin J, and McGuigan M. A brief review of strength and  
329 ballistic assessment methodologies in sport. **Sports Med** 44: 603–623, 2014.
- 330 23. Meylan C, McMaster T, Cronin J, Mohammed I, and Rogers C. Single leg lateral,  
331 horizontal and vertical jump assessment: reliability, interrelationships, and ability to  
332 predict sprint and change of direction performance. **J Strength Cond Res** 23: 1140-  
333 1147, 2009.
- 334 24. Meylan C, Cronin J, Oliver J, Hughes M, and McMaster D. 2012. The reliability of  
335 jump kinematics and kinetics in children of different maturity status. **J Strength**  
336 **Cond Res** 26: 1015-1026, 2012.
- 337 25. Munro A, and Herrington L. Between-session reliability of four hop tests and the  
338 agility t-test. **J Strength Cond Res** 25: 1470-1477, 2011.
- 339 26. Ostojic S, Mazic S, and Dikic N. Profiling in basketball: Physical and physiological  
340 characteristics of elite players. **J Strength Cond Res** 20: 740–744, 2006.
- 341 27. Read P, Oliver J, Myer G, De Ste Croix M, and Lloyd R. The Effects of Maturation  
342 on Measures of Asymmetry During Neuromuscular Control Tests in Elite Male Youth  
343 Soccer Players. **Pediatric Exer Sci** 44: 1-23, 2017.

- 344 28. Reid A, Birmingham B, Starford P, Alcock K. Hop testing provides a reliable and  
345 valid outcome measure during rehabilitation after anterior cruciate ligament injury.  
346 **Phys Ther** 87: 337–349, 2007.
- 347 29. Ross M, Langford B, and Whelan J. Test re-test reliability of 4 single leg horizontal  
348 hop tests. **J Strength Cond Res** 16: 617-622, 2002.
- 349 30. Stolberg M, Sharp A, Comtois A, Lloyd R, Oliver J, and Cronin J. Triple and  
350 Quintuple Hops: Utility, Reliability, Asymmetry, and Relationship to Performance.  
351 **Strength Cond J** 38: 18-25, 2016.
- 352 31. Thomas C, Dos Santos T, Comfort P, and Jones AJ. Between session reliability of  
353 common strength and power- related measures in adolescent athletes. **Sports** 5: 5-15,  
354 2017.
- 355 32. Thomas J, Nelson J, and Silverman S. Research methods in physical activity. 3<sup>rd</sup>  
356 Edition, Champaign, Illinois. Human Kinetics, 2005.
- 357 33. Turner A, Brazier J, Bishop C, Chavda S, Cree J, and Read P. Data analysis for  
358 strength and conditioning coaches: Using excel to analyze reliability, differences, and  
359 relationships. **Strength Cond J** 37: 76-83, 2015.
- 360 34. Young, W. Laboratory strength assessment of athletes. **New Study Athletics** 10: 88–  
361 96, 1995.
- 362 35. Ziv G, and Lidor R. Physical attributes, physiological characteristics, on-court  
363 performances and nutritional strategies of female and male basketball players. **Sports**  
364 **Med** 39: 547–568, 2009.

365

366

367

368

369 Table 1: Standardized warm up protocol followed prior to testing

Phase	Exercise or Drill	Prescription		
Raise heart rate	Light jogging – forward, backwards, side shuffles	5 minutes		
Activate and Mobilize	Mini-band lateral shuffle	1 x 8 each side		
	Floor glute bridges	1 x 8		
	Quadruped thoracic spine extension	1 x 6 each side		
	Bodyweight squats	1 x 10		
	Multi-planar lunge circuit	1 x 6 each side		
	Single leg squat	1 x 5 each side		
Potentiate	Single leg hop and stick	1 x 2 each side		
	Triple hop (stick final landing)	1 x 2 each side		
	Crossover hop (stick final landing)	1 x 2 each side </tr <tr> <td>Single leg countermovement jump</td> <td>1 x 2 each side</td> </tr>	Single leg countermovement jump	1 x 2 each side
	Single leg countermovement jump	1 x 2 each side		

370

371

372

373

374


375

376

Single Leg Hop

Triple Hop

Crossover Hop



377

378

379

380

381

382

383

384

385 Figure 1: Schematic of horizontal hop tests showing number and direction of hops

386

Table 2: Mean data (cm)  $\pm$  standard deviations (SD) for each trial during test session 1, inclusive of within-session reliability data and asymmetry values (calculated from mean scores).

<b>Test</b>	<b>Trial 1 (SD)</b>	<b>Trial 2 (SD)</b>	<b>Trial 3 (SD)</b>	<b>CV (%)</b>	<b>SEM (cm)</b>	<b>ICC (95% CI)</b>	<b>Mean Scores</b>	<b>Asymmetry (%)</b>
SLH (R)	114 $\pm$ 18.7	114 $\pm$ 17.3	115 $\pm$ 18	4.4	0.33	0.89 (0.79-0.95)	119.2 $\pm$ 17.4	1.16 $\pm$ 1.80
SLH (L)	116 $\pm$ 13.7	115 $\pm$ 15.3	117 $\pm$ 14.4	3.8	0.37	0.87 (0.74-0.94)	120.6 $\pm$ 14.8	
TH (R)	367 $\pm$ 56.9	370 $\pm$ 49.2	364 $\pm$ 48.2	3.0	0.25	0.94 (0.88-0.97)	377.4 $\pm$ 52.1	0.24 $\pm$ 3.32
TH (L)	360 $\pm$ 48.4	364 $\pm$ 46.1	372 $\pm$ 46.1	3.6	0.33	0.89 (0.79-0.95)	378.3 $\pm$ 47.4	
CH (R)	298 $\pm$ 54.3	299 $\pm$ 43.9	306 $\pm$ 53	6.0	0.42	0.82 (0.67-0.92)	319.6 $\pm$ 54.4	2.02 $\pm$ 3.54
CH (L)	308 $\pm$ 55.8	315 $\pm$ 61.3	315 $\pm$ 62.9	4.3	0.26	0.93 (0.86-0.92)	326.2 $\pm$ 59.4	
SLCMJ (R)	9.5 $\pm$ 2.6	9.5 $\pm$ 2.4	9.6 $\pm$ 2.5	2.6	0.11	0.99 (0.98-0.99)	9.8 $\pm$ 2.6	5.10 $\pm$ 0.14
SLCMJ (L)	9.2 $\pm$ 2.8	9.0 $\pm$ 2.8	9.0 $\pm$ 2.7	3.0	0.11	0.99 (0.97-0.99)	9.3 $\pm$ 2.8	

R = Right leg, L = left leg, CV = Coefficient of variation, SEM = Standard error of the measurement, ICC = Intraclass correlation coefficient, CI = confidence intervals, SA = Symmetry angle, SLH = Single leg hop, TH = Triple hop, CH = Crossover hop, SLCMJ = Single leg countermovement jump

Table 3: Mean data (cm)  $\pm$  standard deviations (SD) for each trial during test session 2, inclusive of within-session reliability data and asymmetry values (calculated from mean scores).

<b>Test</b>	<b>Trial 1 (SD)</b>	<b>Trial 2 (SD)</b>	<b>Trial 3 (SD)</b>	<b>CV (%)</b>	<b>SEM (cm)</b>	<b>ICC (95% CI)</b>	<b>Mean Scores</b>	<b>Asymmetry (%)</b>
SLH (R)	118 $\pm$ 15.2	118 $\pm$ 13.4	116 $\pm$ 15.4	3.9	0.35	0.88 (0.76-0.95)	121.5 $\pm$ 14.3	0.82 $\pm$ 2.12
SLH (L)	116 $\pm$ 13.5	117 $\pm$ 11.4	115 $\pm$ 12.8	4.1	0.44	0.81 (0.64-0.91)	120.5 $\pm$ 11.3	
TH (R)	376 $\pm$ 42.5	372 $\pm$ 46.6	367 $\pm$ 40.8	3.6	0.35	0.88 (0.76-0.95)	381.4 $\pm$ 45.5	0.81 $\pm$ 2.62
TH (L)	366 $\pm$ 51.2	366 $\pm$ 47.4	370 $\pm$ 44.0	3.3	0.49	0.92 (0.83-0.96)	378.3 $\pm$ 49.2	
CH (R)	308 $\pm$ 40.5	315 $\pm$ 47.1	315 $\pm$ 52.0	5.4	0.23	0.83 (0.69-0.93)	329.4 $\pm$ 44.6	0.33 $\pm$ 4.45
CH (L)	316 $\pm$ 54.4	315 $\pm$ 57.3	319 $\pm$ 54.4	3.6	0.19	0.94 (0.88-0.98)	328.3 $\pm$ 50.9	
SLCMJ (R)	9.5 $\pm$ 2.6	9.5 $\pm$ 2.5	9.6 $\pm$ 2.4	2.8	0.27	0.99 (0.97-0.99)	9.8 $\pm$ 2.5	6.12 $\pm$ 0.14
SLCMJ (L)	9.1 $\pm$ 2.7	8.8 $\pm$ 2.7	8.9 $\pm$ 2.8	3.5	0.16	0.99 (0.97-0.99)	9.2 $\pm$ 2.7	

R = Right leg, L = left leg, CV = Coefficient of variation, SEM = Standard error of the measurement, ICC = Intraclass correlation coefficient, CI = confidence intervals, SA = Symmetry angle, SLH = Single leg hop, TH = Triple hop, CH = Crossover hop, SLCMJ = Single leg countermovement jump

Table 4: Best scores for test sessions 1 and 2  $\pm$  standard deviations (SD), pooled coefficient of variation (CV) data, between-session intraclass correlation coefficient (ICC) data, and the smallest worthwhile change (SWC).

Test	Best Scores (SD)		Pooled CV (%)	ICC (95% CI)	SWC (%)
	Session 1	Session 2			
SLH (R)	119.2 $\pm$ 17.4	121.5 $\pm$ 14.3	4.1	0.76 (0.48-0.89)	2.63
SLH (L)	120.6 $\pm$ 14.8	119.4 $\pm$ 13.4	4.0	0.72 (0.40-0.88)	2.35
TH (R)	377.4 $\pm$ 52.1	381.4 $\pm$ 45.5	3.3	0.87 (0.71-0.95)	2.57
TH (L)	378.3 $\pm$ 45.0	378.3 $\pm$ 49.2	3.5	0.85 (0.65-0.94)	2.49
CH (R)	319.6 $\pm$ 54.4	329.4 $\pm$ 44.6	5.7	0.79 (0.54-0.91)	3.05
CH (L)	326.2 $\pm$ 60.4	328.3 $\pm$ 56.9	4.0	0.84 (0.64-0.94)	3.58
SLCMJ (R)	9.8 $\pm$ 2.6	9.8 $\pm$ 2.5	2.7	0.99 (0.98-0.99)	5.20
SLCMJ (L)	9.3 $\pm$ 2.8	9.2 $\pm$ 2.7	3.2	0.99 (0.98-0.99)	5.95

R = Right leg, L = left leg, SLH = Single leg hop, TH = Triple hop, CH = Crossover hop, SLCMJ = Single leg countermovement jump