

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Cockburn, Emma, Hills, Frank, Wilson, Laura J., Paice, Katherine, Wood, Alyssa and Dimitriou, Lygeri (2017) The effect of plasma volume on changes in biomarkers of muscle damage, inflammation, oxidative stress and leukocyte differential following a 42.2 km trail run, and the efficacy of recovery interventions. In: BASES Conference 2017, 28-29 Nov 2017, Nottingham, UK.

Final accepted version (with author's formatting)

This version is available at: <http://eprints.mdx.ac.uk/23227/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

The effect of plasma volume on changes in markers of muscle damage, inflammation and oxidative stress following a 42.2 km trail run, and the efficacy of recovery interventions

Plasma Volume changes following exhaustive exercise: Does this affect the magnitude of change induced by exercise and/or the interaction effect of recovery interventions?

Cockburn, E., Hills, F., Wilson, L., Paice, K., Wood, A. and Dimitriou, L.

Middlesex University (Middlesex, UK)

Introduction

The use of cold water immersion and/or the consumption of a Montmorency cherry juice to facilitate exercise recovery are becoming increasingly popular. The efficiency of these exercise recovery interventions has been examined by measuring specific circulating biomarkers. However, exercise, can produce a transient fluid shift out of the intravascular space, resulting in haemoconcentration (Alis et al, 2015). The aim of this study was to examine whether plasma volume affects a) the magnitude of change over time in biomarkers and b) the outcomes of these recovery strategies following a 42.2 km trail run.

Methods

Twenty seven endurance trained males (mean age 40 ± 8 years; height 1.77 ± 0.08 m; mass 75.9 ± 10.3 kg) completed a 42.2 km trail run. Participants were randomly assigned to one of 3 recovery interventions (placebo (PL); cherry juice (CJ); or cold water immersion (CWI)). Markers of inflammation (C-reactive protein (CRP); Interleukin-6 (IL-6)), muscle damage (Creatine Kinase (CK)) and oxidative stress (Protein carbonyls (PC); Thiobarbituric acid reactive substances (TBARS); Superoxide dismutase (SOD)) and full blood count were measured before and immediately, 24h and 48h following the run. All biomarkers were adjusted for any plasma volume (PV) changes using the Dill and Costill (1974) equation. The interaction effect between recovery intervention and time at each level of plasma volume (uncorrected (U) and corrected (C)) were analysed using a $3 \times 4 \times 2$ mixed model analyses of variance. Simple main effects analyses were calculated for significant interaction effects between treatment and time by PV. The level of significance was set at $P < 0.05$.

Results

Analysis revealed significant interaction effects of time x PV ($P < 0.05$) for all variables except TNF- α ($P > 0.05$). For CRP and TBARS post-hoc tests revealed significant changes over time for U and C data. There was no significant change over time for SOD U or C. U IL-6 and PC analysis revealed no significant change over time, however there was a significant change over time for C values. Similarly, CK analysis demonstrated significant changes over time for U values but no significant change over time for C values. There were no significant interaction effects of time x PV x group ($P > 0.05$).

Conclusion

A marathon results in significant changes in markers of inflammation (CRP; IL-6), muscle damage (CK) and oxidative stress (PC; TBARS) over 48 hours. Plasma volume affects the magnitude of

change over time in these markers. However, plasma volume did not affect the outcomes of these recovery interventions.

References

Alis, R., Sanchis-Gomar, F., Primo-Carrau, C., Lozano-Calve, S., Dipalo, M., Aloe, R., Blesa, J.R., Romagnoli, M. and Lippi, G. (2015) Hemoconcentration induced by exercise: Revisiting the Dill and Costill equation. *Scandinavian Journal of Medicine and Science in Sports*.

Dill, D.B. and Costill, D.L. (1974) Calculation of percentage changes in volumes of blood plasma, and red cells in dehydration. *Journal of Applied Physiology*.