

Middlesex University Research Repository

An open access repository of
Middlesex University research

<http://eprints.mdx.ac.uk>

Purchase, Diane ORCID: <https://orcid.org/0000-0001-8071-4385>, Bisschop, Lieselot, Ekberg, Christian, Fedotov, Petr, Garelick, Hemda ORCID: <https://orcid.org/0000-0003-4568-2300>, Kandile, Nadia G., Luque, Rafael, Popoola, Oluseun, Rüdell, Heniz, Serpe, Angela, Surati, Kirankumar and Wilson, Benjamin P. (2018) Global approaches to environmental exposure - assessment of e-wastes. In: SETAC Europe 28th Annual Meeting, 13-17 May 2018, Rome, Italy. . [Conference or Workshop Item]

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/24671/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

Global approaches to environmental exposure – assessment of e-waste

Diane Purchase¹, Lieselot Bisschop², Christian Ekberg³, Petr Fedotov⁴, Hemda Garelick¹, Nadia G. Kandile⁵, Rafael Luque⁶, Oluseun Popoola⁷, Heinz Rüdell⁸, Angela Serpe⁹, Kirankumar Surati¹⁰, Benjamin P. Wilson¹¹

INTRODUCTION & OBJECTIVES

The waste stream of obsolete electronic equipment grows exponentially, creating a worldwide problem.

e-waste issues are complex, multi-faceted and can only be successfully tackled via a multidisciplinary, trans-boundary approach that involves all stakeholders that include amongst others: manufacturers, scientists, economists, policy makers, waste professionals and consumers.

The e-waste project aims to highlight:

- i) discrepancies in the provision and enforcement of regulations between developed and emerging countries;
- ii) complexity in the analysis of e-waste contaminants in environmental and biological samples; and
- iii) A harmonised approach should be taken to use compound-specific trace analysis



METHODOLOGY

- A systematic literature review on four key aspects: a) chemical analysis; b) environmental and health impacts; c) recycling and treatment of e-waste and d) governance. The review using databases such as: ISI Proceedings, JSTOR Search, Scopus, Web of Science, PubMed
- Case study analysis from many different countries

RESULTS & DISCUSSION

The e-waste stream comprises of a heterogeneous mix of different metals, metalloid, glass, plastics, flame retardants and valuable materials such as gold, silver, copper and aluminium (Fig. 1). A sizeable amount of e-waste has been discarded in developing countries due to the high cost of safe recycling processes and stringent regulations in developed countries (Fig. 2).

In developed countries, e-waste management revolved around two major strategies: recycling or disposal in their own countries, or exportation to other countries (developed and developing).

In developing countries, the management of e-waste is rudimentary (e.g. dismantling, chipping, melting and burning are often used by the informal sector to recover valuable materials from different e-waste components) and hazardous (Table 1, Fig. 2). These unofficial recycling practices contribute to the release of toxic metals and persistent pollutants that affect both the environment and human health. It is further complicated by illegal import of the waste, hindered by lack of technology, undermined by weak environmental regulations and constrained by inadequate organizational structure.




Fig. 1: Material constituent of some equipment that end up as e-waste

Table 1. Examples of informal recycling activities around the world.

Activities	Country										
	China	Egypt	Ghana	India	Indonesia	Nigeria	Pakistan	Philippines	South Africa	Uruguay	Vietnam
Physical dismantling using bare hands and simple hand tools such as hammers, chisels screw drivers to separate different material	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Removing components from printed circuit boards by burning	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Printed circuit boards sold to informal exporters for further treatment	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Stripping of metals in acid baths	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Crushing and/or melting plastics	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Burning cables to recover precious metals	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Manual stripping of wires	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Burning unwanted materials in open air	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Disposing unsalvageable materials in fields/landfill/riverbanks	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Refilling of toner cartridges	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

Over the last decade, a large body of knowledge has emerged on the risk posed to the environment in e-waste recycling sites (Table 2). It is apparent that there is a lack of harmonisation of tools or indices to assess risk in environment and health, particularly in soil.

Default risk assessment methods frequently **overestimate** exposure by assuming that a chemical will be equally bioavailable in all media, irrespective of the properties of the environmental media or the chemical form of the contaminant.

Using total pollutant concentration in health risk models like to overestimate the potential daily intake of contaminants from informal e-waste recycling sites. Whereas, bioavailability and bioaccessibility data provide more accurate assessment of the risk.

Table 2. Criteria used in literature to assess risk in e-waste recycling sites.

Criteria	Examples
Pollutant's total concentration	Heavy metals in surface dust in India ²
Considering the bioavailability of the pollutants	Heavy metals in soils from Nigeria ³ , dechlorane plus in soil from China ⁴ , heavy metals in soil from Ghana ⁵
Considering the bioaccessibility (physiological based extraction test)	Pb in Philippines ⁶ , heavy metals in China ^{7,8}
Compare with Regulatory benchmarks and control sites	Heavy metals in groundwater ⁹ and well water ¹⁰ in China, soil in India ¹¹
Using pollution indices	PAHs in soil in China ¹² , PCB and BFRs in dust in Vietnam ¹³ , heavy metals in soil in China ¹⁴ and Ghana ¹⁵
Using risk models to examine exposure pathways	Dioxin/PCDD/Fs in soil ¹⁶ and heavy metals in air in China ^{17,18}

PAHs = Polycyclic aromatic hydrocarbons; PCB = Polychlorinated biphenyl; BFRs = Brominated flame retardants; PCDD/F = Polychlorinated dibenzo-p-dioxins and dibenzofurans

CONCLUSION

- i) Significant discrepancies exist in the provision and enforcement of regulations between developed and emerging countries.
- ii) The analysis of e-waste contaminants in environmental and biological samples is complex and challenging.
- iii) Conceptual models should be used to ascertain the source-pathways-receptors route of the pollutants
- iv) Currently, there is lack of harmonisation of tools or indices to assess risk in environment and health, particularly in soil.
- v) It is proposed that a harmonised approach should be taken to use appropriate speciation analysis (e.g. to assess bioavailable or bioaccessible fractions) to evaluate e-waste contaminant risk.



Fig. 2: Transboundary routes of e-waste transport¹



Fig. 3: Informal recycling activities in developing countries

ACKNOWLEDGEMENTS

The e-waste project "*The Environmental and Health Challenges of e-waste and its Management: an Emerging 21st Century Global Concern*" (#2014-031-3-600) is supported by the International Union of Pure and Applied Chemistry (IUPAC). The project brings together multidisciplinary global expertise to explore different aspects of the e-waste challenge: chemical analysis of contaminants, policy and governance, environmental and health impacts, development and advances in treatment technologies including e-waste valorisation.

Email: d.purchase@mdx.ac.uk
IUPAC project webpage: https://iupac.org/projects/project-details/?project_nr=2014-031-3-600

Addresses:

¹Department of Natural Sciences, Faculty of Science and Technology, Middlesex University, London NW4 4BT, UK
²Department of Criminology, Erasmus University Rotterdam, Burgemeester Oudlaan 50, Rotterdam, the Netherlands
³Division of Energy and Materials, Department of Chemical and Chemical Engineering, Chalmers University of Technology, Kemivägen 4, S-41296 Gothenburg, Sweden
⁴Russian Academy of Sciences, Vernadskii Institute, 19 Kosygin St, RF-119991, Moscow
⁵Department of Chemistry, Faculty of Women, Ain Shams University, Heliopolis, Cairo, Egypt
⁶Departamento de Química Orgánica, Universidad de Córdoba, Rabanales Campus, Marie Curie Building, National Court IV-A, km. 396.14014-Córdoba, Spain
⁷Department of Chemical Science, Yaba College of Technology, Yaba, Lagos, Nigeria.
⁸Fraunhofer Institute for Molecular Biology and Applied Ecology (Fraunhofer IME), 57392 Schmallenberg, Germany
⁹Dipartimento di Ingegneria Civile, Ambientale e Architettura, Università degli Studi di Cagliari, Via Marengo, 2, Edificio ex DIGITA Cagliari, Italy
¹⁰Department of Chemistry, Sardar Patel University, Vallabh Vidyanagar- 388120, Anand,Gujarat, India
¹¹Department of Chemical and Metallurgical Engineering, School of Chemical Engineering, Aalto University, P.O. Box 16200, 00076 Aalto, Finland

References

<http://www.sustainelectronics.illinois.edu/policy/international.cfm> [accessed 7/9/16]
¹Awasthi AK et al. 2016. <https://doi.org/10.1016/j.envpol.2015.11.027>
²Isimekhai KA et al. 2017. <https://doi.org/10.1007/s11356-017-8877-9>
³Li L et al. 2014. [https://doi.org/10.1016/S1001-0742\(13\)60447-7](https://doi.org/10.1016/S1001-0742(13)60447-7)
⁴Tokumaru T et al. 2017. <https://doi.org/10.1007/s00244-017-0434-5>
⁵Fujimori T et al. 2017. <https://doi.org/10.1016/j.ihazmat.2017.07.066>
⁶Fao XQ et al. 2014. <https://doi.org/10.1007/s11356-014-3562-8>
⁷Han Z et al. 2017. <https://doi.org/10.1080/09064710.2016.1222916>
⁸Zheng J et al. 2013. <https://doi.org/10.1016/j.ecoenv.2013.06.017>
⁹Wu Q et al. 2015. <https://doi.org/10.1016/j.scitotenv.2014.10.121>
¹⁰Pradham JK & Kumar S. 2014. <https://doi.org/10.1007/s11356-014-2713-2>
¹¹Wang Y et al. 2012. <https://doi.org/10.1016/j.scitotenv.2012.08.018>
¹²Tue NM et al. 2013. <https://doi.org/10.1016/j.envint.2012.11.006>
¹³Zhao W et al. 2015. <https://doi.org/10.1007/s10646-015-1532-7>
¹⁴Kyere VN et al. 2017. doi: [10.5620/ehp.e2016006](https://doi.org/10.5620/ehp.e2016006)
¹⁵Man YB et al. 2015. <https://doi.org/10.1007/s11356-014-3909-1>
¹⁶Zheng X et al. 2016. DOI: [10.4209/aahr.2014.11.0292](https://doi.org/10.4209/aahr.2014.11.0292)
¹⁷Huang C-L et al. 2016. <https://doi.org/10.1016/j.ihazmat.2016.05.081>