

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Skinner, T. ORCID: <https://orcid.org/0000-0002-0018-6963>, Byrne, M., Dickinson, J. K., Fisher, L. ORCID: <https://orcid.org/0000-0001-9481-9727>, Funnell, M., Guzman, S., Hendrieckx, C., Hermanns, N. ORCID: <https://orcid.org/0000-0002-9610-0975>, Kanc, K., Lloyd, C. ORCID: <https://orcid.org/0000-0002-8863-3069>, Mocan, A., Nouwen, Arie ORCID: <https://orcid.org/0000-0002-0609-4082>, Pouwer, F., SalehStattin, N., Snoek, F., Speight, J. ORCID: <https://orcid.org/0000-0002-1204-6896>, Sturt, J., Vallis, M. ORCID: <https://orcid.org/0000-0002-0165-5936>, Wagner, J., Willaing, I., YoungHyman, D. and Zoffmann, V. (2019) Comment on the consensus report on the management of hyperglycaemia in Type 2 diabetes by the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetic Medicine*, 36 (7) . pp. 911-912. ISSN 1464-5491 (doi:10.1111/dme.13934)

Final accepted version (with author's formatting)

This version is available at: <http://eprints.mdx.ac.uk/26379/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

Title: Comment on the Consensus Report on the Management of Hyperglycemia in Type 2 Diabetes by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD).

We read with interest the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD) joint Consensus Report on the Management of Hyperglycemia in Type 2 Diabetes [1]. Following the publication of the report, discussions of the content among networks of behavioural and educational researchers prompted us to write this letter

This consensus report is to be commended for recommending that “providers and health care systems should prioritize the delivery of patient-centered care”. The report is also to be commended for providing some clear actions that can be taken to promote person-centred care, including emphasizing the importance of diabetes self-management education and support, promoting shared decision-making between the person with diabetes and the health care professional, and pointing to some methods that may be beneficial in achieving this. At the same time, we also appreciate that there are challenges in implementing a person-centred approach and we would like to highlight and comment on some of these challenges we see in the Consensus Report.

The first is that the report uses the term “adherence” throughout, making the recommendation that “Facilitating medication adherence should be specifically considered when selecting glucose-lowering medications”. Whilst we agree that it is important to discuss the issues surrounding an individual’s willingness and ability to take any prescribed medications, we suggest that the word “adherence” is used ill-advisedly in the report. A recent series of position statements from Diabetes Australia [2], jointly by the ADA and American Association of Diabetes Educators [3] and jointly by Diabetes UK and the UK’s National Health Service [4], have pointed to the negative consequences of using the terms “adherence” and “compliance”, when talking with or about people living with diabetes. These position statements echo earlier commentaries that argue that compliance and adherence are dysfunctional concepts and empirical work demonstrating that the assumptions underpinning these words are not substantiated [5]. The continued use of the term “adherence”, which implies an unequal relationship in which people with diabetes passively follow clinicians’ instructions, is at best unhelpful. The fact that diabetes care is largely stuck in this “adherence” paradigm may explain why a series of Cochrane reviews on interventions to promote adherence

have repeatedly concluded that “Current methods of improving medication adherence for chronic health problems are mostly complex and not very effective” [6], and argued that the problem needs to be reconceptualised. This can be done readily by active adoption of a “strengths-based, empowering language” [3] and other recommendations of the above-mentioned position statements [2-4] to facilitate active engagement in self-care, health and well-being.

We are also concerned with the statement that “effective consultations include motivational interviewing”. “Motivational interviewing” is a specific set of intervention strategies and techniques requiring extensive training and significant time to deliver. Consequently, it is not possible for most health professionals to include “motivational interviewing” in routine clinical and education practice. Furthermore, recommendations for integrating a specific set of tools and techniques into clinical practice require a high-quality evidence base. However, this does not exist for motivational interviewing in diabetes or chronic disease care [7]. Whilst meta-analyses indicate a possible small benefit of such interventions, this is conflated with the additional time investment. We suggest that the emphasis needs to be placed not on specific methods but on the principles underpinning “motivational interviewing”, common to many effective psycho-educational interventions and essential to effective diabetes care. These principles move the clinician away from an “adherence” paradigm, to focus not on persuading the person with diabetes to take a certain medication, but rather to raise awareness of the individual’s choices and enable agreement about which management option best fits the individual’s needs, goals, values and preferences.

As part of the goal to individualise care, the Consensus Report also provides guidance on the assessment of “key patient characteristics”; specifically, recommending assessment of “Issues such as motivation and depression”. This seems at odds with ADA’s 2016 position statement on the psychosocial care of people with diabetes, which recommends considering an “assessment of symptoms of diabetes distress, depression, anxiety, and disordered eating and of cognitive capacities” [8], and the ADA’s 2019 standards of diabetes care [9]. Thus, we are surprised that this joint Consensus Report does not specifically recommend assessment of diabetes distress, especially given a recent meta-analysis, which indicates that about one third of individuals with type 2 diabetes experience clinically significant levels of diabetes distress [10]. We are also surprised that the report recommends the assessment of “motivation”, given that this term is vague, often used inappropriately, and is not easily assessed. We acknowledge that achieving a paradigm shift takes time, and requires persistent, consistent and insistent

advocacy. However, it is disappointing that this joint Consensus Report does not appear to be implementing the recent recommendations of the ADA, and other international bodies, to incorporate strengths-based language and related strategies, with the aim of enabling widespread promotion of truly person-centred diabetes care. Therefore, we would like to recommend that future similar endeavours (consensus reports, guidelines, position statements) include representation of a broader range of disciplines and relevant associations and study groups e.g. American Association of Diabetes Educators (AADE), Psycho-Social Aspects of Diabetes Study (PSAD) group of the EASD, and the U.S. based Behavioral Research In Diabetes Group Exchange (BRIDGE).

Funding: None

Conflicts of interest: The authors declare no conflicts of interest

Disclaimer: The opinions expressed in this letter reflect those of the authors and not of their institutional affiliations.

Authors:

Timothy Skinner 1,2, Molly Byrne 3, Jane K. Dickinson 4, Larry Fisher 5, Martha Funnell 6, Susan Guzman 7, Christel Hendrieckx 8, Norbert Hermanns 9, Karin Kanc 10, Cathy Lloyd 11, Andreia Mocan 12, Arie Nouwen 13, Frans Pouwer 14, Nouha Saleh-Stattin 15, Frank Snoek 16, Jane Speight 8,17, Jackie Sturt 18, Michael Vallis 19, Julie Wagner 20, Ingrid Willaing 2, Deborah Young-Hyman 21, Vibeke Zoffmann 1.

Affiliations:

1. University of Copenhagen, Denmark
2. Steno Diabetes Centre Copenhagen, Denmark
3. National University of Ireland, Ireland
4. Teachers College Columbia University, U.S.A.
5. University of California San Francisco, U.S.A.
6. University of Michigan Medical School, U.S.A.
7. Behavioural Diabetes Institute, U.S.A.
8. Australian Centre for Behavioural Research in Diabetes, Diabetes Victoria, Australia
9. Jazindiabetes, (Diabetes & Me), Slovenia
10. Forschungsinstitut Diabetes-Akademie Bad Mergentheim (FIDAM GmbH)
11. The Open University, U.K.
12. Emergency Clinical County Hospital, Cluj-Napoca, Rumania
13. Middlesex University, U.K.
14. University of Southern Denmark, Denmark
15. Academic Primary Health Care Centre, Stockholm, Sweden
16. Amsterdam UMC, Netherlands
17. Deakin University, Australia
18. Kings College London, U.K.

19. Dalhousie University, Canada
20. UConn Health, U.S.A.
21. National Institute of Health, U.S.A.

Corresponding author:

Timothy Skinner
Københavns Universitet
Institut for Psykologi
Center for Sundhed og Samfund
Københavns Universitet
Øster Farimagsgade 2A,
1353 København K

Tel: +45 35 33 60 82

Email: ts@psy.ku.dk

ORCID ID: [0000-0002-0018-6963](https://orcid.org/0000-0002-0018-6963)

Word count: 870

Tables / Figures: 0

References

1. Davies MJ, D'Alessio DA, Fradkin J, Kernan WN, Mathieu C, Mingrone G, Rossing P, Tsapas A, Wexler DJ, Buse JB. Management of Hyperglycemia in Type 2 Diabetes, 2018. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care*. 2018 Dec;41(12):2669-2701.
2. Speight J, Conn J, Dunning T, Skinner TC on behalf of Diabetes Australia (2012). A new language for diabetes: improving communications with and about people with diabetes. *Diabetes Research and Clinical Practice*. 97(3) 425-431.
3. Dickinson JK, Guzman SJ, Maryniuk MD, O'Brian CA, Kadohiro JK, Jackson RA, D'Hondt N, Montgomery B, Close KL and Funnell MM. The Use of Language in Diabetes Care and Education. *Diabetes Care* 2017; 40(12): 1790-1799.
4. Cooper A, Kanumilli N, Hill J, Holt RIG, Howarth D, Lloyd CE, Kar P, Nagi D, Naik S, Nash J, Nelson H, Owen K, Swindell B, Walker R, Whicher C, Wilmot E. Language matters. Addressing the use of language in the care of people with diabetes: position statement of the English Advisory Group. *Diabet Med*. 2018 Dec;35(12):1630-1634.
5. Glasgow RE, Anderson RM. In diabetes care, moving from compliance to adherence is not enough. Something entirely different is needed. *Diabetes Care*. 1999 Dec;22(12):2090-2.

6. Nieuwlaat R, Wilczynski N, Navarro T, Hobson N, Jeffery R, Keepanasseril A, Agoritsas T, Mistry N, Iorio A, Jack S, Sivaramalingam B, Iserman E, Mustafa RA, Jedraszewski D, Cotoi C, Haynes RB. Interventions for enhancing medication adherence. *Cochrane Database Syst Rev.* 2014 Nov 20;(11):CD000011.
7. Frost H, Campbell P, Maxwell M, O'Carroll RE, Dombrowski SU, Williams B, Cheyne H, Coles E and Pollock A. Effectiveness of Motivational Interviewing on Adult Behaviour Change in Health and Social Care Settings: A Systematic Review of Reviews. *PLoS ONE* 13(10): e0204890
8. Young-Hyman D, de Groot M, Hill-Briggs F, Gonzalez JS, Hood K and Peyrot M. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. *Diabetes Care* 2016; 39(12): 2126–2140.
9. Lifestyle Management: *Standards of Medical Care in Diabetes-2019*. American Diabetes Association. *Diabetes Care.* 2019 Jan;42(Suppl 1):S46-S60.
10. Perrin NE, Davies MJ, Robertson N, Snoek FJ and Khunti K. The Prevalence of Diabetes-specific Emotional Distress in People with Type 2 diabetes: a Systematic Review and Meta-analysis. *Diabetic Medicine* 2017; 34(11): 1508-20.