

# Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Virgile, Adam and Bishop, Chris ORCID: <https://orcid.org/0000-0002-1505-1287> (2021) A narrative review of limb dominance: task specificity and the importance of fitness testing. *Journal of Strength and Conditioning Research*, 35 (3) . pp. 846-858. ISSN 1064-8011 [Article] (doi:10.1519/JSC.0000000000003851)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/30854/>

## Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

[eprints@mdx.ac.uk](mailto:eprints@mdx.ac.uk)

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

# A Narrative Review of Limb Dominance: Task-Specificity and the Importance of Fitness Testing

## Abstract

Preferential limb function must be sustained through repetitious asymmetrical activities for continuous athletic development and ultimately, optimal athletic performance. As such, the prevalence of limb dominance and between-limb differences are common in athletes. Severe between-limb differences have been associated with reductions in athletic performance and increased injury risk in athletes. However, in the current literature, the terms limb preference and limb dominance have been used inter-changeably. Together, these terms include a limb which is subjectively preferred and one that is objectively dominant in one or more performance measures from a variety of athletic tasks. In this review, we 1) discuss reported correspondence between task-specific limb preference and limb dominance outcomes in athletes, 2) provide greater context and distinction between the terms *limb preference* and *limb dominance*, and 3) to offer pragmatic strategies for practitioners to assess context-specific limb dominance. A limb which is subjectively preferred is not necessarily objectively dominant in one or more athletic qualities or sport-specific tasks. Further to this, a limb which is objectively superior in one task may not exhibit such superiority in a separate task. Thus, limb preference and limb dominance are both task-specific. As such, we propose that practitioners intentionally select tasks for limb dominance assessment which resemble the most relevant demands of sport. Because limb dominance profiles are inconsistent, we suggest that practitioners increase assessment frequency by integrating limb dominance testing into standard training activities. This will allow practitioners to better understand when changes reflect sport-specific adaptation versus potential performance or injury ramifications.

**Key Words:** Limb preference; consistency; between-limb differences; fitness testing

## Introduction

Human limb preference appears to manifest itself during infancy, but the development of limb preference is highly malleable (47, 151). Repeated motor task exposure impacts the degree of limb preference; a range of motor tasks, both complex and simple, involving tools and other objects (e.g., unilateral teeth-brushing, shooting a basketball or throwing a baseball) can contribute to limb preferences (151). Consequently, it is not uncommon for limb preference to differ by body segment. For example, approximately 90% of people exhibit a well-defined right-hand preference, whilst only 25-45% demonstrate right leg preference in lower extremity actions (52). Furthermore, limb preference is evidently task-specific (76, 178). Velotta et al. (178) asked healthy college-aged individuals about limb preference when performing different tasks; 90% of the cohort preferred the right limb to kick a ball, but only 40% preferred the right limb for single-leg standing (178). Thus, it seems important to make the distinction between subjectively preferred and objectively better-performing limbs. Historically, a “dominant” limb has been used in both contexts (e.g., a subjectively preferred limb and an objectively better-performing limb) (168, 175). Further to this, it is not uncommon to see these terms used in study design and analyses without reporting how the “limb dominance” or “limb preference” was determined (73, 74, 118, 180). Choosing to analyze between-limb differences by a subjectively preferred limb (e.g., limb preference) can result in a completely different interpretation than by objective performance (e.g., limb dominance). For example, Kuki et al. (108) found no inter-limb difference in peak force generation when limb performance was defined by preference, but one limb generated significantly greater peak force than the other when limb performance was segregated by dominance (108). The ramifications of not having a clear, well-recognized distinction between limb preference and dominance are clear: researchers cannot accurately consolidate findings relating to asymmetry and limb preference-dominance interactions, and clinicians cannot effectively interpret and use findings in practice. In this review, *limb preference* indicates the subjectively preferred limb for completing a task, whereas *limb dominance* indicates the limb which objectively outperforms the other in a particular task. A relationship can exist between these two constructs — the subjectively preferred limb may also produce better objective performance abilities (3) — but, this is not always the case (103).

Compared with the general population, the interaction between limb preference and limb dominance is more complex in athletes. Sporting tasks oftentimes encompass a tapestry of orchestrated movements, which can be impacted by numerous factors, including sport-specific environmental constraints, spatial orientation of the body, fatigue, and injury (42, 49, 81, 83, 116, 128, 153). In many sports, sustained preferential function through repetitious asymmetrical activities may be required for continuous athletic development and ultimate success (41, 94-97, 101, 152, 161). Athletes may also develop limb preference for sport-specific tasks (115, 131, 164), which may differ from the preferred limb in everyday activity (89, 115). Through asymmetric task-specific repetition, task-specific limb dominance can emerge, or preceding limb dominance can become more pronounced. Associations between limb dominance and reductions in physical performance (6, 11, 21, 23, 32, 36, 95, 114, 119, 120, 122, 123, 150) and increased injury risk (10, 50, 58, 82, 98, 144, 157, 162, 173) have been identified in athlete populations. For example, greater isometric midthigh pull (IMTP) peak force asymmetry was associated with reduced vertical jump height during squat jump and countermovement jump tasks, in collegiate athletes (6). In National Hockey League (NHL) players, preseason hip strength was compared with subsequent in-season adductor strain probabilities (173). Players who sustained adductor strains during the season had lower hip adduction-to-abduction strength ratios in the injured limb compared with the non-injured limb during preseason testing (173). However, there are also investigations where associations between limb dominance and performance or injury risk are not observed (33, 43, 58, 113). Thus, because limb dominance can have potential physical performance and injury implications, its quantification and systematic assessment is likely to be of interest to both practitioners and athletes.

In sport, limb dominance is task-dependent (22, 119, 125, 176). Referring to a limb as being holistically “dominant” from performance in one task is an over-simplification and lacks necessary context. For example, Dos’Santos et al. (59) found that the limb that performed best during horizontal jumping did not correspond with the better-performing limb during change-of-direction (CODS) tasks, in collegiate athletes (59). In soccer, the limb that is used to strike the ball is generally considered dominant (29, 92, 144), but this notion fails to appreciate that the kicking motion is complex and requires the integration of multiple skills (24). Specifically, the limb used to strike a ball may be capable

of producing more force or achieve faster movement velocity, but the standing (e.g., non-kicking) limb may be considered dominant when performing stability or supporting actions (8, 45, 130, 150, 161). Not only should test selection for limb dominance profile assessment reflect the most frequent and relevant demands of sport, but also, test results should be contextualized. Because limb dominance profiles can also change throughout a season of sport participation (18), it is also important to understand when changes in these tasks reflect sport-specific adaptation versus potential performance or injury ramifications.

Therefore, the primary aim of this review is to discuss reported correspondence between task-specific limb preference and limb dominance outcomes in athletes. Secondary aims include: 1) to provide greater context and distinction between the terms *limb preference* and *limb dominance* so as to offer practitioners with a more consistent approach in their understanding and analysis, and 2) to offer pragmatic strategies for coaches and practitioners to assess context-specific limb dominance.

### **Limb Preference vs. Limb Dominance: A Need for Terminology Clarification**

In the current literature, many tasks have been used to determine limb preference in athletes, including the preferred kicking or jumping leg (12, 27, 70, 85, 99, 129, 132, 137, 152), writing, throwing, or serving hand (4, 44, 68, 91, 176), and other various multi-component assessments (e.g., the Lateral Preference Inventory and Waterloo Footedness Questionnaire) (14, 71, 78, 89, 171, 174). The heterogeneity and generalized methodology used to quantify limb preference in athletes is confusing and like limb dominance, limb preference in one task cannot be inferred from its preference in another (178). The Lateral Preference Inventory (LPI) is a questionnaire frequently used to categorize limb preference (48). In the LPI, people are asked about which leg they would prefer to use during bilateral mobilizing tasks (e.g., tasks involving both legs), including: 1) “With which foot would you kick a ball to hit a target?”, 2) “Which foot would you use to step on a bug?” and, 3) “If you had to step up onto a chair, which foot would you place on the chair first?” (48). In each of these tasks, the non-participating limb maintains contact with the ground (e.g., when stepping onto a chair, one foot is planted on the ground to support the body while the other foot steps onto the chair). Hart and Gabbard (77) categorized limb preference of 100 university students using the LPI (77). Students

were also asked to identify the limb preferred to perform a single-leg balance task whereby the other limb was not in contact with the ground. More than half of students (62%) who had right limb preference via LPI categorization (e.g., bilateral context) preferred the left limb for performing the static single-leg balance task (e.g., unilateral context). Almost half of students (44%) who had left limb preference on the LPI also switched limb preference for the static single-leg balance task (77). In other words, the task used in limb preference assessment influenced the choice of limb preferred. While “With which foot would you kick a ball to hit a target?” is on the LPI and has contextual appropriateness for a soccer player, it would be an unsuitable question to infer unilateral jumping preference for a basketball or volleyball athlete (71, 133, 160). Thus, in order to gain meaningful insights from limb preference assessment in athletic performance contexts, the task in which the limb is preferred must be carefully considered. An array of tasks and metrics have also been used to quantify limb dominance, including medial knee displacement during bilateral jumping (130), muscle strength (104, 166, 169), balance (27, 110, 182), unilateral jumping (105, 132, 152), and change-of-direction performance (97), among various other assessments (42, 55, 122). This review will focus on interactions between limb preference and limb dominance for balance, muscle strength, unilateral jumping, and change-of-direction task performances because they are often assessed in literature and demonstrate contextual application for many sports.

### **Understanding Sport-specific Tasks**

Athletic success in sport oftentimes requires repetitious high-level performance of specific biomechanical patterns. For actionable limb preference and dominance appraisal, performing a needs analysis for the sport in question is necessary and will provide practitioners with the information they need to choose assessments that align with the sport-specific biomechanical demands. To categorize motor tasks in sport, Maloney (122) suggests using the motor task groupings presented by Guiard (90): (Group 1) unilateral (e.g., throwing a baseball), (Group 2) bilateral asymmetric (e.g., hockey shot), (Group 3) out-of-phase bilateral symmetric (e.g., sprinting or cycling), and (Group 4) in-phase bilateral symmetric (e.g., bilateral jump or weightlifting). Because sport can encompass a wide array of motor tasks, holistic sport categorization is difficult, and it is arguably more relevant to categorize

the primary tasks required in the sport of interest. For example, batting in baseball (127) requires bimanual asymmetric high-velocity actions (Group 2), whereas throwing (69) requires unimanual high-velocity actions (Group 1). Batting in cricket (28) or using the stick in ice hockey (140) represent other bimanual asymmetric high-velocity actions, whereas throwing in handball, shot put (111), or hammer represent other unimanual high-velocity actions. As such, because a repetitious unimanual movement is required to throw a baseball, limb dominance of high magnitude in a throwing arm should be expected for this action, and the benefit of limb dominance appraisal may not outweigh the time, resource, and energy costs to routinely assess its presence. For example, unimanual overhead athletes oftentimes present with relatively greater internal rotation strength than external rotation strength (2, 54, 56, 65, 67, 148, 179) in the preferred (e.g., throwing or serving) shoulder, compared with the non-preferred shoulder. Thus, it may not be important or realistic for an athlete to spend time aiming to restore this between-limb imbalance.

The range of motor tasks and associated groupings that warrant assessment are sport-dependent; each sport sits on a continuum in respect to both relevance and frequency of motor task and motor task group requirement. Most sports, such as soccer and basketball, rely on frequent high-level performance of Group 2 and Group 3 tasks. Although other task groups are relevant (e.g., layups: Group 1; bilateral jumping: Group 4), kicking, approach jumping, and changing direction (Group 2), as well as sprinting (Group 3), are particularly relevant in these sports. Because soccer and basketball rely on frequent high-level performance of tasks in Group 2 and Group 3 categories, two-sided proficiency is likely required to enhance athletic success probability (7, 34, 39, 88, 164, 165). Through frame-by-frame video analysis, Carey et al. (40) analyzed foot use patterns in a sample of 236 soccer players from 16 teams in the 1998 World by comparing passing, first touch, dribble and tackle success rates between preferred and non-preferred limbs (40). Players demonstrated bias towards preferred foot use during games, but the skill levels (e.g., success rates) between preferred and non-preferred limbs were similar (40). Players recognize the significance of limb dominance as it pertains to skill acquisition. When asked about what it takes to be a skilled soccer player, 400 amateur soccer players determined that “two-footedness” (e.g., being equally skilled with both feet) was a very important quality (39). Stöckel and Vater (164, 165) reported similar results in professional, semi-

professional, and amateur basketball players. Frame-by-frame video analysis of 14 games revealed a bias towards using the preferred hand for all assessed skills (dribbling, passing, catching, and throwing) (165). A negative linear relationship ( $r = -0.39$ ) between preferred hand use and level of competitive play was reported; the frequency of ball contacts with the preferred hand decreased from 59.2% in amateurs to 49.6% in semi-professionals to 48.8% in professional players (165). Like soccer players, elite basketball players recognize the importance of equal skill acquisition across limbs; when questioned about hand preference for performing basketball-specific tasks, nearly all players ( $n=176$ ) reported that being equally skilled with both hands was necessary for basketball success (164).

As described above, whether limb preference and dominance presence will benefit athletic performance depends on many variables, including the sport, task, and context. Because some sports require bimanual and bipedal task-specific proficiency for success (e.g., dribbling in soccer and basketball), devoting substantial efforts toward enhancing abilities of the non-preferred or non-dominant limb may be advantageous, but such efforts may not be warranted in other sports or tasks. Understanding the demands of the sport and its most relevant tasks through a needs analysis is a prerequisite for contextually suitable limb preference and limb dominance profiling. If assessed strategically, longitudinal profiling can potentially enhance athletic performance through improved training prescription and a better understanding of limb dominance underpinnings.

### **Limb Preference and Limb Dominance: Do they Correspond?**

Balance, various muscle strength measurements, unilateral jumping and change-of-direction performance appear to be the most common tasks used for limb dominance assessments in literature, and thus, will form the focus of this review. Just because a limb is preferred does not infer that it will objectively outperform the non-preferred limb in balance (76), strength (108), unilateral jumping (131), or change-of-direction tasks (72). In the following sections, the correspondence between limb preference and limb dominance for these tasks will be discussed in detail.



### ***Balance Tasks***

Bressel et al. (27) assessed static and dynamic balance performances using the Balance Error Scoring System (BESS) (145) and the Star Excursion Balance Test (SEBT) (87), respectively, in a group of 34 National Collegiate Athletic Association (NCAA) Division I female athletes (soccer, n = 11; basketball, n = 11; gymnastics, n = 12), (27). There were no meaningful differences in the SEBT or BESS performances between the non-preferred and preferred (e.g., preferred limb to kick a ball) limbs, in any of the sport groups (27). The modified SEBT is also termed the Lower Quarter Y Balance Test (YBT-LQ) and includes three reaching directions: anterior, posteromedial, and posterolateral. In an investigation of YBT-LQ performance, Fort-Vanmeerhaeghe et al. (72) quantified limb preference in elite youth female basketball players by asking players to identify which limb was preferred to: 1) kick a ball, 2) initiate stair climbing, and 3) regain balance following a slight, unexpected perturbation, with the limb chosen in two or more of the scenarios considered as the preferred limb. There was little consistency between limb preference and dominance; only 45%, 48% and 41% players demonstrated better objective YBT-LQ performance on the preferred limb in the anterior, posteromedial, and posterolateral directions, respectively (72).

Comparable limb preference and dominance discrepancies are also observed in rugby athletes. Brown et al. (31) reported inconsistent limb preference-dominance results in male rugby union athletes during a dynamic evaluated dynamic single leg balance task during more stable and less stable conditions, using a Biodex Balance SD system (31). The Biodex Balance SD system measures the degree of tilt about each axis, and it is from this assessment that anterior-posterior, medial-lateral, and overall stability indices were quantified and used to determine balance performance (31). Limb preference was defined by the preferred kicking leg. Rugby union backs performed moderately worse on the non-preferred limb for medial-lateral and overall stability indices in the more stable (Biodex Balance Level 8) condition, but there were no differences between non-preferred and preferred limbs in the anterior-posterior stability index or any of the less stable (Biodex Balance Level 2) stability indices (31). In addition, there were no differences between non-preferred and preferred limbs in any of the stability indices for the forwards (31). Static and dynamic balance performances were assessed in 37 elite Australian Footballers using force plate excursion metrics and

the results were similar; there were no differences between non-preferred and preferred (e.g., preferred to kick a ball) limbs in static or dynamic balance performances (102). In contrast, limb preference coincided with limb dominance in a large cohort of skiers. Steidl-Müller et al. (159) investigated balance performances in 285 high-level competitive ski athletes (125 females, 160 males) from three different age categories: 95 youth (aged 10–14 years), 107 adolescent (aged 15–19 years), and 83 elite athletes (aged 20–34 years) (159). Athletes were asked to balance on an MFT® Challenge Disc for 20 seconds on each limb, with the level of stability (e.g., stability index) recorded based on the position of the body's center of gravity within the circle. Youth, adolescent, and elite skiers attained  $10.9 \pm 9.5\%$ ,  $11.3 \pm 7.7\%$ , and  $9.6 \pm 6.8\%$  worse stability index scores on the non-preferred limb, compared with the preferred (e.g., preferred to accept body weight upon being instructed to “fall forward”) limb (159).

Using the Upper Quarter Y Balance Test (YBT-UQ), Borms et al. (26) found no differences in YBT-UQ performance between non-preferred and preferred (e.g., preferred to throw a ball) limbs in 29 overhead athletes (26). Ludwig et al. (117) reported larger dynamic knee valgus angle during single-leg drop landing down from a box in the preferred (e.g., preferred to kick a ball) limb, in 114 amateur and elite youth soccer players (117). Further to this, elite players had larger between-limb differences than their amateur counterparts. The kinematic differences between limbs reflect the different motor requirements of the supporting and the kicking leg in soccer. The kinematic differences between cohorts reflect a potential impact of sport-specific adaptation on limb preference-dominance relationships.

The cohorts assessed along with limb preference and limb dominance determination methods in the balance literature are heterogeneous. With this consideration, the available body of evidence suggests that the preferred limb rarely coincides with a performance advantage in static or dynamic balance tasks. There are many potential reasons as to why these discrepancies exist. For example, the method used for limb preference determination may have been unsuitable in that it did not include similar biomechanical demands to the tasks used for limb dominance assessment. The limb preference criteria often included mobilizing tasks (e.g., limb used to kick a ball or initiate stair climbing) while the limb dominance task assessed was static balance. The disparate speed and force of movement

between preferred and dominance criteria may also be a factor (e.g., kicking a ball is a forceful action requiring high velocity of movement, while SEBT or YBT-LQ reaching is a low-force and low-velocity activity). Further to this, kicking a soccer ball is a bipedal activity (one foot kicks the ball while the other supports the body), while many of the outcome measures required unilateral stance. Peters (138) sums up this notion by stating that “in the design of [lower extremity limb] controls, the preferred limb should perform the actions that are more directly related to the goal of movement,” (138). To make relevant comparisons between limb preference and dominance outcomes, motor task group categories and important biomechanical patterns should be similar between limb preference and dominance assessments. Interactions between cohort demographics, including injury history, age, training age, access to training and recovery resources, and sport-specific adaptations, among various other factors, will also likely influence these limb preference-dominance relationships.

### ***Muscle Strength Tasks***

Isokinetic dynamometry is the most common muscle strength assessment in limb preference literature to date. Although relatively novel in the literature, functional field test assessments using force plate and motion capture technology is becoming more prevalent in practice. Thus, these two methods of muscle strength assessment will be the focus of the literature discussed below.

Rahnama et al. (141) investigated differences between non-preferred and preferred (e.g., preferred to kick a ball) limbs in concentric and eccentric isokinetic knee flexion and extension strength (peak torque) at varying speeds (1.05, 2.09, and 5.23 rad/s) in 41 elite and sub-elite English male soccer players aged  $23.4 \pm 3.8$  years (141). No differences were found in any of the muscle strength measures between the non-preferred and preferred limbs except for the knee flexors at 2.09 rad/s during concentric muscle actions, where the non-preferred limb demonstrated greater strength than the preferred limb ( $119 \pm 22$  vs.  $126 \pm 24$  Nm) (141). Daneshjoo et al. (53) reported similar findings in a slightly younger cohort of 36 male professional soccer players, aged  $18.9 \pm 1.4$  years. There were no differences in isokinetic knee flexion peak torque, knee extension peak torque, or hamstring:quadricep (H:Q) peak torque ratios at varying speeds (60, 180, and 300 deg/s) between non-preferred and preferred (e.g., preferred to kick a ball) limbs (53). In 17 female national team

soccer players (aged  $24.2 \pm 3.7$  years), Maly et al. (124) also found no differences between non-preferred and preferred (e.g., preferred to kick a ball) limbs in H:Q peak torque ratios at varying speeds, including 60, 120, 180, 240, and 300 deg/s (124). Similar to older cohorts, Maly et al. (125) also found no differences between non-preferred and preferred (e.g., preferred to kick a ball) limbs in isokinetic H:Q peak torque ratios at 60, 120, 180, 240, or 300 deg/s in 41 male under-16 national team soccer players (125). More successful soccer players demonstrate a high H:Q peak torque ratio, particularly at high isokinetic velocity. The H:Q ratio tends to increase in both limbs as velocity increases (124), and this phenomenon is thought to occur due to greater antagonistic activation of the hamstring muscles with increasing isokinetic velocity, leading to a subsequent increase of the H:Q ratio (167). In elite soccer players, the H:Q ratio may increase more for the non-preferred limb than the preferred limb (168). This notion has also been reported in world-class tennis and squash athletes. Read et al. (142) reported no difference between non-preferred and preferred (e.g., the leg ipsilateral to the racket arm) limbs for isokinetic H:Q peak torque ratio at 90 deg/s, but at faster speeds (180 deg/s, 240 deg/s, and 300 deg/s), greater H:Q ratios for the non-preferred limb were reported (142). However, a disproportional increase in H:Q ratio for the non-preferred limb with increasing velocity is not always observed (66). The notion that the hamstrings produce more force compared to the quadriceps as speed of movement increases, particularly in the non-preferred leg (9, 142, 147, 170), indicates that a complex relationship exists between lower limb musculature and single joint movement velocity, which may be influenced by many factors, including: neural demands, muscle architecture, and mechanical advantage. According to a limited quantity of research in basketball players, relationships between the preferred limb and isokinetic dynamometry metrics seem to echo those of the aforementioned cohorts. No significant differences were evident between non-preferred and preferred (e.g., preferred to kick a ball) limbs in isokinetic knee flexion or knee extension peak (60 and 180 deg/s) in 12 professional basketball players (167).

In contrast, other studies compare outputs between dominant (e.g., objectively better-performing) and non-dominant (e.g., objectively worse-performing) limbs (59, 104, 105, 109), and this is an important distinction when interpreting between-limb differences (108). Kuki et al. (108) investigated bilateral and unilateral isometric midhigh pull (IMTP) capabilities in 15 male collegiate

football players and sprinters. There were no differences between the non-preferred and preferred (e.g., preferred to kick a ball) limbs in bilateral or unilateral IMTP peak force generation (108). Interestingly, more than half of athletes performed better with the non-preferred limb; during bilateral and unilateral IMTP tasks, 10/15 (67%) and 8/15 (53%) athletes, respectively, generated greater peak force with the non-preferred limb compared with the preferred limb (108). Furthermore, when peak force was compared between non-dominant (e.g., less peak force) and dominant (e.g., greater peak force) limbs, statistically significant inter-limb differences were observed during both bilateral ( $p < 0.05$ ; Cohen's  $d = 1.12$ ) and unilateral ( $p < 0.05$ ; Cohen's  $d = 0.56$ ) IMTP tasks (108). During the bilateral IMTP, muscle activity (mean EMG as a percentage of unilateral muscle activity) for the gluteus maximus, gluteus minimus, semitendinosus, biceps femoris, rectus femoris, and vastus lateralis muscles were also collected. There were no differences in muscle activity for any muscle groups between non-preferred and preferred limbs, but when segregated by limb dominance (e.g., the limb with higher EMG values compared to the limb with lower EMG values), the dominant limb had significantly higher vastus lateralis muscle activity than the non-dominant limb (108). In another study by Kuki et al. (109), between-limb comparisons for unilateral IMTP, countermovement jump, and drop jump performances were assessed in 20 male collegiate football players (109). Six performance metrics were assessed (IMTP absolute peak force, IMTP relative peak force, IMTP absolute peak force at 100ms, IMTP relative peak force at 100ms, countermovement jump height, and reactive strength index), and the non-preferred limb was outperformed by the preferred (e.g., preferred to kick a ball) limb in only one of the metrics: IMTP relative peak force (109). Gleason (79) assessed between-limb bilateral IMTP peak force in 17 NCAA Division I male soccer players. Significant differences between non-preferred and preferred (e.g., preferred to kick a ball) limbs were evident, and only 9/17 (53%) of athletes produced greater peak force with the preferred limb (79). Kobayashi et al. (106) investigated inter-limb ground reaction force differences in male long jumpers while performing the back squat exercise with loads of 50, 70, and 90% of their three-repetition maximum (106). There were no differences in peak vertical and horizontal ground reaction forces between non-preferred and preferred (e.g., leg used for long jump takeoff) limbs in any loading condition. However, bilateral kinematic differences, particularly at the hip joint, were reported (106). While

athletes may not present with preference-related strength imbalances, limb preference may influence the adoption of asymmetric movement strategies to achieve bilaterally equivalent strength levels.

Isokinetic and IMTP muscle strength limb dominance cannot be inferred from identifying the limb that is subjectively preferred using traditional limb preference assessment methods. When considered relevant for athletic success, practitioners are encouraged to assess muscle strength limb dominance directly. Kinematic analyses may provide additional value to identify disparate movement strategies across limbs, particularly when kinetic outcomes are similar. Regardless of the sport-specific demands of the athletic cohort assessed, limb preference was quantified as the limb that was preferred to kick a ball (in most studies), which may have influenced limb preference-dominance correspondence outcomes. Strength assessment was constrained to mostly non-functional tasks by way of isokinetic dynamometry; isokinetic assessments typically have poor relationships with functional athletic performance tasks, such as sprinting and jumping (51, 135). More research investigating limb preference-dominance interactions in muscle strength tasks is warranted. Future research should investigate correspondence between functional field tests and limb preference for performance of a similar motor task in order to better understand how limb preference practically relates to limb dominance in sport.

### ***Jump, Change-of-Direction Speed (CODS), and Sprint Tasks***

Many sports require high-level proficiency in a tapestry of specific power movements. As such, these movements are arguably the most sought-after quality in sport and are frequently tested as physical performance measures (60, 62, 63, 155). Because jumping, CODS, and sprinting are common to most sports and are the most studied power-related tasks in the limb preference literature, limb preference-dominance relationships for these tasks will be discussed below.

Vaisman et al. (174) studied limb preference-dominance interactions in 27 professional soccer players (174). To determine limb preference, players were instructed to: 1) kick a soccer ball, 2) extinguish a simulated fire, and 3) draw figures on the ground; the limb that was mobilized in two or more of the three tasks was determined to be preferred. There were no differences in unilateral vertical jump height or power between preferred and non-preferred limbs (174). Samadi et al. (149)

conducted a study comparing limb preference and unilateral single-leg horizontal triple jump performance in one-legged athletes (e.g., long and high jumpers) and two-legged athletes (no description given). There were no differences in jumping performances between non-preferred and preferred (e.g., preferred to kick a ball) limbs, in either group (149). Mulrey et al. (132) examined relationships between limb preferences and unilateral vertical and horizontal jumping performances in 40 adolescent (aged  $15.5 \pm 1.2$  years) female basketball players (132). Limb preference was categorized into kicking and jumping preferences. Players were asked to respond to two different questions: 1) “What limb would you use to kick a ball as far as possible?” and, 2) “What limb would you use to jump as high as possible?”, determining kicking and jumping limb preference, respectively. The preferred limb for jumping did not outperform the non-preferred jumping limb in the unilateral vertical (single-leg countermovement jump) or horizontal (single-leg triple hop jump for distance) jumping tasks. The preferred limb for kicking performed better than the non-preferred kicking limb in the single-leg triple hop for distance task, but not the single-leg countermovement jump task (132). Fort-Vanmeerhaeghe et al. (71) assessed unilateral countermovement jump performance to examine limb preference-dominance correspondence in 79 volleyball and basketball athletes (41 males, 38 females) (72). To determine limb preference, athletes were asked to identify the preferred limb with which to: 1) kick a ball, 2) initiate stair climbing and, 3) regain balance following a slight and unexpected perturbation. The limb that was identified in two or more of the three responses was determined to be preferred. Less than half (40.5%) of athletes achieved greater countermovement jump height when jumping with the preferred limb (71). Using the same limb preference criteria as the previous study, Fort-Vanmeerhaeghe et al. (72) assessed inter-limb differences in multidirectional unilateral jumping (vertical, horizontal, and lateral directions) and change of direction speed ([CODS]; 10 meter total distance with a  $180^\circ$  turn at 5 meters) performances in 29 elite youth female basketball players (72). Similar to previous findings, less than half of players demonstrated objectively better unilateral jumping performance on the preferred limb; only 48%, 48% and 35% of players performed better with the preferred limb during unilateral vertical, horizontal, and lateral jump tasks, respectively. On the CODS task, roughly half of players (52%) achieved faster times when the preferred limb was also the plant limb for the  $180^\circ$  turn (72). Greska et al. (85) reported no between-

limb differences when evaluating hip and knee biomechanics between non-preferred and preferred (e.g., preferred to kick a ball) limbs during an unanticipated side-cutting task in 20 collegiate female soccer athletes (85). Joint angles, internal moments, vertical ground reaction forces at initial contact, peak knee adductor moment and peak stance periods did not differ between limbs upon planting, despite equivalent approach velocities. Additionally, there were no differences in muscle activity (peak EMG as a percentage of maximal voluntary isometric contraction) at any of the time points assessed for any of the measured muscles, including the gluteus medius, biceps femoris, semitendinosus, rectus femoris, vastus lateralis, and vastus medialis (85).

Brown et al. (30) investigated force and power outputs during initial acceleration and maximal velocity sprinting in 30 male academy-level rugby union players (30). The forwards produced lower peak relative vertical force, peak relative horizontal force, and maximal relative power with the non-preferred limb than the preferred (e.g., the “front” leg during sprint setup) limb during acceleration (effect size [ES] = -0.32, -0.58 and -0.67, respectively) and maximal velocity sprinting (ES = -0.50, -0.65 and -0.60, respectively). During acceleration, the backs produced similar between-limb vertical forces (ES = 0.02) with unclear differences in horizontal forces and maximal power. The backs also produced similar vertical forces (ES = 0.10) between limbs during maximal sprinting, alongside an unclear between-limb difference in maximal power and greater horizontal force (ES = 0.54) with the non-preferred limb (30). Korhonen et al. (107) investigated between-limb force outputs during sprinting in 18 younger (age  $23 \pm 4$  years) and 25 older (aged  $70 \pm 4$  years) high-level sprinters and found no differences in mean force outputs between the non-preferred and preferred (e.g., preferred to perform a single-leg jump) limbs (107).

When considering the available evidence, there does not appear to be a consistent correspondence between the limb which is subjectively preferred and the limb which is objectively dominant for jumping, CODS, and sprinting performances in athletes. Although correspondence between limb preference and limb dominance during balance, muscle strength, and power performance tasks may exist in athletes, evidence dissemination is obstructed by a host of difficulties: 1) few studies on the topic, 2) heterogeneity in methods and cohorts assessed, 3) heterogeneity in limb



preference quantification criteria, and 4) lack of limb preference criteria specificity as it relates to the task performance of interest.

### **Limb Dominance: A Comparison Between Sports-specific Tasks**

Because sporting success requires high-level performance of complex motor actions that include varying combinations of kinematic, biomechanical, and kinetic properties, it is not uncommon for a limb to objectively outperform the other in one sporting action, but not another. For example, a limb that is dominant in strength tasks may not be dominant in jumping or CODS tasks. Dos'Santos et al. (59) investigated correspondence between horizontal jumping and CODS abilities in 22 male collegiate team sport athletes (soccer, n =10; rugby, n = 6; cricket, n = 6) and found that the better-performing (e.g., dominant) limb for horizontal jumping did not correspond to the plant limb which produced faster times for 180° and 90° turns during CODS (59). In another study by Dos'Santos et al. (57), unilateral IMTP force-time characteristics, including relative peak force and impulse at 200 and 300ms, were compared with between-direction CODS performance in 20 male collegiate athletes (soccer: n=8, rugby: n=6, and cricket: n=6) (57). The dominant limb for IMTP metrics was rarely dominant for CODS performance (via the 505 test); nearly half of athletes (9/20) had significant between-limb IMTP relative peak force differences, and of the nine athletes who had significant strength asymmetry, only one (11%) produced faster times when the stronger limb was also the plant limb for 180° turns during CODS. Similarly, 6/20 and 7/20 athletes had significant between-limb IMTP impulse differences at 200 and 300ms, respectively, but only one athlete had a significantly faster CODS time when planting on the limb that also had greater IMTP impulse values (57). Gleason (79) compared IMTP strength with CODS times and ground foot contact times in 17 NCAA Division I soccer athletes; unilateral IMTP limb strength was determined through bilateral IMTP performance on dual force plates and the 505 test was used to quantify CODS performance (79). The limb which was considered dominant for peak force during the IMTP task was not the same limb which was considered to be dominant in the 505 CODS task (79).

When comparing performances between two different jumping tasks, a limb that is dominant in one task may not also demonstrate dominance in the other. Using a portable force platform, Bishop

et al. (16) compared unilateral isometric squat (peak force), countermovement jump (peak force, concentric impulse, and eccentric impulse), and broad jump (peak force, concentric impulse, and eccentric impulse) performances in 28 recreational sport athletes (16) and reported task-dependent limb dominance (16). Calculating a Kappa coefficient is an approach used to assess the consistency of a between-limb difference (e.g., limb dominance) for a common metric across two tasks and describes the proportion of agreement between two methods after any agreement by chance has been removed (16, 46). When assessing limb dominance consistency in peak force, Kappa values ranged from poor to slight (-0.34 to 0.05); limb dominance peak force consistency was slight between the unilateral isometric squat and countermovement jumps (Kappa = 0.04), poor between the isometric squat and broad jumps (Kappa = -0.34), and slight between the countermovement and broad jumps (Kappa = 0.05) (16). Five of the six descriptive levels of agreement between the dominant and non-dominant limbs for eccentric and concentric impulse metrics during the two jumping tasks ranged from poor to fair (Kappa = < 0.01 to 0.32), with only concentric impulse between unilateral countermovement and broad jumps showing substantial levels of agreement (Kappa = 0.79) (16). In a separate study, Bishop et al. (17) investigated limb dominance profiles of under-17 elite female soccer players, again using a portable force platform (17). Players performed unilateral squat jumps, countermovement jumps, and drop jumps; jump height, peak force, concentric impulse, and peak power were reported as common metrics across tasks. Again, inconsistent limb dominance was present across tasks. Kappa coefficients revealed fair to substantial levels of agreement for direction of limb dominance between the unilateral squat and countermovement jump tasks (Kappa = 0.35 to 0.61), but only poor to fair levels of agreement between the squat jump and drop jump (Kappa = -0.26 to 0.18) and countermovement and drop jump tasks (Kappa = -0.13 to 0.26) (17). During a mid-season fitness test battery inclusive of unilateral countermovement, lateral, and broad jump tasks, Madruga-Parera et al. (120) evaluated between-limb differences in 42 elite youth handball athletes (120). Similar to findings of Bishop et al. (17, 22), the limb that was dominant in one task was rarely dominant in another, as demonstrated by Kappa coefficients ranging from poor to slight (-0.05 to 0.15) (120); differences between unilateral countermovement and lateral jumps (Kappa = 0.15), countermovement and broad jumps (Kappa = -

0.05), and lateral and broad jumps (Kappa = 0.00), were slight, poor, and poor agreements, respectively (120).

Limb dominance for one task or athletic quality (e.g., strength, power, and agility) cannot be used to infer limb dominance in another. An athlete that achieves greater peak force with one limb during IMTP may not be more agile when pivoting on that limb during a CODS task or achieve greater height with that limb during vertical jumping. Even when tasks have many similar qualities (e.g., unilateral vertical, horizontal, and lateral jumping), it cannot be inferred that a better-performing limb in one task will perform better in the other, similar tasks. The task-specific nature of limb dominance is important for practitioners to understand when assessing inter-limb qualities to make more informed decisions for athlete care and development. It is suggested that practitioners consider the athletic qualities of the task being assessed when applying data in practice from limb dominance and asymmetry evaluation.

### **Limb Dominance: A Task-specific Concept**

The notion that limb dominance is task-specific is well-recognized throughout the literature (8, 22, 45, 72, 79, 95, 120, 141, 156) and can be further articulated by understanding specific movement patterns that occur in sport. To better contextualize this, we provide an example of the interaction between offensive and defensive actions required in the sport of basketball.

When a player has to perform a skill when interacting with other players (e.g., in games and team practices) or under high pressure circumstances, the most proficient limb in the given situation will be most often used to perform the skill (163). Using data on 3,647 National Basketball Association (NBA) players from 1946 through 2009, Lawler et al. (112) discovered that the vast majority of NBA players prefer to shoot with the right hand; the overall prevalence of right-hand preference (e.g., preferred hand to shoot the ball) of NBA players was 94.9% (112). Thus, during NBA competition, most players will use the right hand for layup attempts. A right-handed layup attempt includes a 2-step approach sequence whereby the left leg ultimately propels the athlete into the air, creating the required vertical displacement to perform the action successfully (158). Within the layup skill itself, the two-step sequence involves a horizontal displacement focus during the first

step to create separation from the defender, and a vertical displacement on the second step to bring the athlete closer to the basket for increased scoring success (92). Sensibly, the repetitious favoring of one side in a unilateral task, such as a layup, is thought to increase the likelihood of an athlete developing a lower-body muscular imbalance (158), because one lower limb is consistently tasked with facilitating maximal center of mass displacement in the horizontal direction while the other limb is tasked with doing the same, but vertically. Therefore, right-hand layup preference may facilitate horizontal and vertical displacement dominance development for the left and right lower limbs, respectively. A general NBA right-hand favoritism for offensive actions, including layups, may also have an impact on limb dominance development in NBA players on the defensive side of the ball. In response to right-hand offensive preference, defenders are required to repetitiously perform actions that may develop lateral horizontal displacement dominance in the right lower limb. On defense, a basketball player must consistently perform explosive lateral movements while facing the offensive player in attempt to prohibit the offensive player from getting close to the basket with the ball (126). In fact, defensive shuffling makes up the largest proportion of high-intensity activity during elite basketball competition (3.1%), followed by sprinting (2.8%), striding (2.4%), sideways running (1.9%), and jumping (1.3%) (1). The defensive player is tasked with prohibiting the offensive player from doing what he or she prefers, which includes using the right side for performing common offensive skills, including dribbling and accelerating towards the basket for a layup opportunity. To prevent the offensive player from executing skills with the preferred (e.g., right) side, the defensive player must perform more frequent and explosive lateral horizontal displacement actions with the right lower limb. A basketball player who performs a higher proportion of layup attempts from the right side on offense (requiring vertical center of mass displacement facilitated by the left limb), and a higher explosive lateral horizontal actions with the right lower limb on defense to prevent their right-handed counterparts from utilizing the preferred limb, may create a task-specific limb dominance profile whereby the left and right lower limbs exhibit vertical and lateral horizontal jump dominance, respectively. Using the aforementioned example, the prevalence of task-specific limb dominance should be expected in basketball athletes. Monitoring task-specific limb dominance changes over time would allow for practitioners to better understand the relevance of such changes and whether they are

due to sport-specific adaptations. Given that inconsistencies in limb dominance over repeated test sessions and time points appears to be evident (18,19) repeated testing seems warranted in order to build a consistent picture of limb dominance characteristics, before such data can be used to inform decision-making.

### **Limb Dominance: Importance of Longitudinal Monitoring**

Limb dominance testing using tasks which reflect the demands of an athlete's sport is of high importance. First, the requirements of the athlete to tolerate the demands of his or her position in sport must be determined by conducting a needs analysis. There is a substantial body of literature covering the demands of various sports (5, 15, 61, 136, 143, 172); however, it is imperative that a current needs analysis is conducted due to the time-sensitive changes in participating athletic populations, equipment, and sport rules, that take place. Once the kinetic, kinematic, and physiological qualities required for successful sport performance are well understood, the practitioner has the knowledge to develop tests which resemble the athletic qualities of interest. It is through the informed selection of tests, each intentionally chosen to provide insight into a specific feature of successful sport performance, that relevant task-specific limb dominance appraisal can occur.

The timing and frequency of limb dominance testing, as well as the method applied to calculate the metrics used for determining limb performance, are important considerations. Because directional limb dominance between test sessions can be inconsistent (20), frequent limb dominance profile assessment is likely necessary to differentiate between consistent limb dominance and fluctuations in natural performance variability. Using a portable force plate, Bishop et al. (20) assessed agreement between unilateral isometric squat (peak force and impulse), countermovement jump (peak force and jump height), and drop jump (reactive strength index [RSI] and jump height) performances across two testing sessions separated by three days, in 28 recreational soccer and rugby athletes (20). Between-session consistency of limb dominance direction for the unilateral isometric squat, countermovement jump, and drop jump, were fair to substantial ( $Kappa = 0.29$  to  $0.64$ ), substantial ( $Kappa = 0.64$  to  $0.66$ ), and fair to moderate ( $Kappa = 0.36$  to  $0.56$ ), respectively (20). This data indicates that the direction of asymmetry can be highly variable. Further to this, levels of

agreement were higher for tests with greater reliability; the unilateral countermovement jump test had the lowest coefficient of variation (CV) for both within and between test sessions ( $CV \leq 6.3\%$ ), and also had the highest levels of agreement (Kappa = 0.64 to 0.66) (20). Using the average of testing trials as the metric to calculate inter-limb differences was more consistent across sessions than using the data from the best trial; thus, calculating the average of trials, as opposed to using the best trial for each limb, might be considered more appropriate for assessing inter-limb differences (20). In another study, Bishop et al. (19) assessed athletic performance in 18 elite male under-23 male academy soccer players at three time points over the course of a competitive season: at pre-season, mid-season, and end-season (19). Unilateral countermovement jump (jump height and concentric impulse) and drop jump (jump height and RSI) performances were measured, and magnitude and direction of limb dominance were assessed independently. Limb dominance magnitude changed in a trivial-to-small, nonlinear fashion throughout the season for both unilateral countermovement jump (ES range = -0.43 to 0.05) and drop jump (ES range = -0.18 to 0.41) performances, providing the impression of consistent asymmetry scores over time. However, limb dominance direction was extremely variable. Levels of agreement ranged from poor to substantial for both the unilateral countermovement jump (Kappa = -0.06 to 0.77) and drop jump (Kappa = -0.10 to 0.78) measures. Substantial levels of agreement were only shown for jump height when comparing mid-season with end-season (Kappa = 0.68) and RSI when comparing pre-season with mid-season (Kappa = 0.78); all other time points showed poor to fair levels of agreement for the direction of limb dominance (19). As such, this highlights the need to: 1) test over repeated sessions to identify any consistencies in limb dominance characteristics (or lack thereof), and 2) consider the direction and magnitude independently when interpreting limb dominance outcomes.

Given that levels of agreement for limb dominance appears highly variable, practitioners are encouraged to adopt regular and longitudinal testing protocols to establish an athlete's limb dominance characteristics more accurately. Further to this, frequent limb dominance assessment will allow practitioners to make a distinction between whether inherent changes over time are merely a consequence of natural fluctuations in performance variability, caused by sport-specific adaptation, or illuminate an underpinning injury or performance concern. When testing limb dominance, conducting

multiple trials per test, and using average metric values (rather than peak or “best” values) are recommended for limb dominance calculations.

### **Practical Applications**

Limb dominance in one test does not infer dominance in another. Therefore, when selecting performance tests for limb dominance assessment, practitioners should carefully consider the demands of sport and positional requirements to ensure test and sport demands are similar (100, 168). From a practitioner’s perspective, functional field tests (e.g., jumps, IMTP, CODS) are recommended because they can be more easily integrated into training environments compared with more laboratory-based assessments (e.g., isokinetic dynamometry), which also typically have poor relationships with athletic performance tasks (51, 135).

**\*\* INSERT TABLE 1 ABOUT HERE \*\***

Although coincidence between limb preference and dominance during balance tasks was discussed, balance is not included as a primary limb dominance assessment parameter because it is rarely considered a key physical testing parameter for athlete populations. When surveyed, an overwhelming proportion of strength coaches working in Major League Baseball (MLB) (64), the National Football League (NFL) (62), NHL (63), and NBA (155) reported testing strength, power, and agility qualities as part of physical testing; with the exception of a few strength coaches in the NHL. However, no strength coaches reported testing balance.

Infrequent limb dominance testing during a competitive season may cause mis-interpretation of limb dominance testing results, owing to the frequent fluctuations seen in limb dominance characteristics (18-20). Making important decisions based off a single data point is inherently risky, particularly when the data lacks consistency and agreement. Therefore, more frequent testing (e.g., weekly to monthly) is suggested so that practitioners can better identify trends in limb dominance profiles and prescribe appropriate training, accordingly. By increasing testing frequency, and thus, the

number of testing data points, practitioners will be better able to detect true changes in athletic performance and limb dominance if and when they exist (13).

In sport training environments, and in field testing, there are many factors that can confound results, which may affect their interpretation, and ultimately, effective implementation of longitudinal performance monitoring (84). For example, differing warm-up protocols, prior training load exposures, testing times, and nutrition practices between testing sessions may influence observed performance changes. Thus, it is of utmost importance that practitioners try to control for as many potential confounding variables as possible to facilitate the capture of contextually valid and reliable results. As such, adopting a standardized testing protocol is suggested. Components of this standardized protocol may include a specific warmup sequence prior to testing, instructions given during testing, technologies used (if any), and footwear selection. Other potential contextual performance moderators, including test timing, are important to consider. Selecting a specific day within the training schedule (e.g., morning of game day or the day following a game day-off day sequence), and time of day to implement testing will reduce the factors that could contribute to performance variability, and thus, increase confidence in observed results. There are differences in resistance training performances between morning and evening times, but habitually training at the same time of day is likely to mitigate these differences (25, 86). An additional benefit of selecting a consistent day and time for repeated testing is that it allows for athletes and staffs to develop a routine.

Standardized repeated testing is suggested. However, we recognize that testing standardization that resembles the strict controls of a lab setting may not always be entirely feasible in some sporting environments due to a host of challenges, such as, chaotic training environments, limited staff support, dynamic competition schedules, and limited training time. These real-world challenges make it difficult to routinely collect valid and reliable data. Given this reality, we will discuss strategies that attempt to minimize confounding influence on results that can be pragmatically applied in sport settings with high frequency: 1) using data outputs from technologies already implemented in structured training with sport coaches (e.g., global position system [GPS] and inertial measurement units [IMUs]), and 2) integrating testing as a normal part of the training routine. Effective execution of either of these strategies require considerable forethought; similar to



conducting a research experiment, a plan that is ecologically feasible and well-controlled must be in place prior to implementation.

Collaboration between practitioners, and coaches is advised to enhance sport-specific performance (35, 38, 139). When players are instrumented with a GPS unit or IMUs on each limb during training with coaches (e.g., practice), sport-specific limb dominance data is available. However, collaboration with sport coaches is necessary to foster a recurring controlled environment that will allow for meaningful application data outputs as they relate to limb dominance. To gain longitudinal insights, we suggest developing a consistent testing framework around the coaches' practice philosophies and drill tendencies. For example, the coaches may prefer to run "Drill X" following days off from training. For example, the practitioner and coaches might collectively decide that once per month that "Drill X" is prescribed for five minutes in duration as the second drill in the drill sequence for that day. Spatio-temporal gait metrics can be used to quantify limb dominance, which can be derived from a single GPS unit worn during training (80, 134, 181). When interpreting results, it is important to appreciate the validity and reliability of the metrics and technology, particularly for GPS devices (93, 121, 154, 177). Although a natural consequence of technology evolution, validity, and reliability within and across devices can impact longitudinal limb dominance data interpretation. Another means of limb dominance quantification during sport-specific training may include between-limb IMU ground reaction force comparisons (37, 146). To our knowledge, limb dominance between sport-specific training using GPS or IMUs and functional field tests has not yet been investigated. Because limb dominance is task-specific (8, 22, 45, 95, 156), observations during sport-specific training may not coincide with observations from performances in functional field tests.

Typically, practitioners have greater control over how activities are structured in their training environment. Thus, functional field testing in this environment is an ideal place for practitioners to facilitate testing in an organized and consistent manner, even if standardized warmups and direct practitioner-led testing are infeasible. Prescribing sets of complex training (e.g., a set of loaded exercise followed by a set of plyometric exercise) (75) within a session as part of the program design, may facilitate more frequent limb dominance data collection, assuming that athletes are able to record

performance of the plyometric movement. For example, five sets of a loaded lower body exercise with 3-5 unilateral vertical jumps in between sets would be an example of an exercise pairing using complex training. When training groups are large and technology is used for data collection, athletes may be instructed to only track jumping performance for one of the five sets, allowing for collection of 3-5 data points per athlete without the burden of waiting for technology to become available. During a consistently programmed unilateral loaded exercise, recording weight lifted alongside rating of perceived exertion or repetitions in reserve per limb is a simple, technology-free means for collecting information that could provide limb dominance data from the training environment. The examples above are not suggested implementations. Rather, they should be viewed as ideas that facilitate critical thinking for how to strategically apply more frequent limb dominance testing in a practical manner that is minimally burdensome to practitioners, coaches, and athletes. With forethought and strategic implementation, more frequent limb dominance testing is possible in many sport training environments. If strategies that can be pragmatically applied with high frequency in real-world sport settings do not exist, effective longitudinal limb dominance assessment in these settings cannot occur. Infrequent limb dominance testing, which is known to provide inconsistent results (18-20), would be the alternative approach. Despite the strategies offered lacking the full scientific rigor of standard experiments, high-frequency testing that minimizes burden to athletes coaches using semi-standardized protocols (using as much standardization as possible) will enable practitioners to better understand limb dominance profiles over time, fostering more informed monitoring and program design decisions for athletes.

## **Conclusion**

A limb which is subjectively preferred in one task is not always the same limb which is preferred for a separate task. Furthermore, a subjectively preferred limb does not always exhibit superiority in objective athletic performance, including balance, strength, jumping, or sprinting-based tasks. Not only are there discrepancies between subjectively preferred and objectively dominant limbs within individuals, but also, a limb that is objectively superior in one task may also be objectively inferior in a separate task. In other words, similar to limb preference, limb dominance is task-specific.

Given the task-specific nature of limb dominance, testing tasks which reflect the demands of an athlete's sport is of critical importance for practitioners.

An athlete's direction of limb dominance appears to be inconsistent when assessed infrequently over the course of a competitive season (e.g., pre-season, mid-season, and end-season). Due to the intra-individual specificity of magnitude and direction of limb dominance, inter-limb relationships, whether it be between test metrics or changes over time (i.e. longitudinal), should occur on the individual level. In order to determine whether limb dominance is consistent or merely fluctuations in performance variability, more frequent limb dominance testing is suggested. Incorporating semi-standardized testing within the training environment or using information collected during sport-specific training that occurs under supervision of the sport coaches are strategies that may allow for increased testing with minimal burden to athletes and coaches. In this review, we assigned the preferred limb label to the limb that is subjectively preferred whereas the dominant limb referred to the limb that is objectively dominant in a given task. Future research should aim to clarify distinctions between limb preference and dominance.

## References

1. Abdelkrim NB, Castagna C, Jabri I, Battikh T, El Fazaa S, and El Ati J. Activity profile and physiological requirements of junior elite basketball players in relation to aerobic-anaerobic fitness. *The Journal of Strength & Conditioning Research* 24: 2330-2342, 2010.
2. Alfredson H, Pietilä T, and Lorentzon R. Concentric and eccentric shoulder and elbow muscle strength in female volleyball players and non-active females. *Scandinavian journal of medicine & science in sports* 8: 265-270, 1998.
3. Annett M. The growth of manual preference and speed. *British Journal of Psychology* 61: 545-558, 1970.
4. Astolfi MM, Struminger AH, Royer TD, Kaminski TW, and Swanik CB. Adaptations of the shoulder to overhead throwing in youth athletes. *Journal of Athletic Training* 50: 726-732, 2015.
5. Baechle TR and Earle RW. *Essentials of Strength Training and Conditioning*. Human kinetics, 2008.
6. Bailey C, Sato K, Alexander R, Chiang C-Y, and Stone MH. Isometric force production symmetry and jumping performance in collegiate athletes. *Journal of Trainology* 2: 1-5, 2013.
7. Bale P and Scholes S. Lateral dominance and basketball performance. *Journal of Human Movement Studies* 12: 145-151, 1986.
8. Barone R, Macaluso F, Traina M, Leonardi V, Farina F, and Di Felice V. Soccer players have a better standing balance in nondominant one-legged stance. *Open Access Journal of Sports Medicine* 2: 1, 2011.
9. Baroni BM, Ruas CV, Ribeiro-Alvares JB, and Pinto RS. Hamstring-to-quadriceps torque ratios of professional male soccer players: a systematic review. *The Journal of Strength & Conditioning Research* 34: 281-293, 2020.
10. Belhaj K, Meftah S, Mahir L, Lmidmani F, and Elfatimi A. Isokinetic imbalance of adductor–abductor hip muscles in professional soccer players with chronic adductor-related groin pain. *European journal of sport science* 16: 1226-1231, 2016.
11. Bell DR, Sanfilippo JL, Binkley N, and Heiderscheit BC. Lean mass asymmetry influences force and power asymmetry during jumping in collegiate athletes. *Journal of Strength and Conditioning Research/National Strength & Conditioning Association* 28: 884, 2014.
12. Bencke J, Curtis D, Krogshede C, Jensen LK, Bandholm T, and Zebis MK. Biomechanical evaluation of the side-cutting manoeuvre associated with ACL injury in young female handball players. *Knee Surgery, Sports Traumatology, Arthroscopy* 21: 1876-1881, 2013.
13. Biau DJ, Kernéis S, and Porcher R. Statistics in brief: the importance of sample size in the planning and interpretation of medical research. *Clinical orthopaedics and related research* 466: 2282-2288, 2008.
14. Bini RR, Jacques TC, Sperb CH, Lanferdini FJ, and Vaz MA. Pedal force asymmetries and performance during a 20-km cycling time trial. *Kinesiology: International journal of fundamental and applied kinesiology* 48: 193-199, 2016.
15. Bishop C, Brazier J, Cree J, and Turner A. A needs analysis and testing battery for field hockey. *Professional Strength & Conditioning* 36: 15-26, 2015.
16. Bishop C, Lake J, Loturco I, Papadopoulos K, Turner AN, and Read P. Interlimb asymmetries: The need for an individual approach to data analysis. *Journal of Strength and Conditioning Research*, 2018.
17. Bishop C, Pereira LA, Reis VP, Read P, Turner AN, and Loturco I. Comparing the magnitude and direction of asymmetry during the squat, countermovement and drop jump tests in elite youth female soccer players. *Journal of Sports Sciences*: 1-8, 2019.
18. Bishop C, Read P, Bromley T, Brazier J, Jarvis P, Chavda S, and Turner AN. The association between Inter-limb asymmetry and athletic performance tasks: a season long study in elite academy soccer players. *Journal of Strength and Conditioning Research*, 2019 (in press).

19. Bishop C, Read P, Chavda S, Jarvis P, Brazier J, Bromley T, and Turner A. Magnitude or Direction? Seasonal Variation of Interlimb Asymmetry in Elite Academy Soccer Players. *Journal of Strength and Conditioning Research*, 2020.
20. Bishop C, Read P, Chavda S, Jarvis P, and Turner A. Using unilateral strength, power and reactive strength tests to detect the magnitude and direction of asymmetry: A test-retest design. *Sports* 7: 58, 2019.
21. Bishop C, Read P, McCubbine J, and Turner A. Vertical and Horizontal Asymmetries are Related to Slower Sprinting and Jump Performance in Elite Youth Female Soccer Players. *Journal of Strength and Conditioning Research*, 2018.
22. Bishop C, Turner A, Lake JP, Read P, Loturco I, and Papadopoulos K. Inter-limb asymmetries: The need for an individual approach to data analysis. *Journal of Strength and Conditioning Research* 40: 1-6, 2018.
23. Bishop C, Turner A, and Read P. Effects of inter-limb asymmetries on physical and sports performance: a systematic review. *Journal of Sports Sciences* 36: 1135-1144, 2018.
24. Bishop DT, Wright MJ, Jackson RC, and Abernethy B. Neural bases for anticipation skill in soccer: an fMRI study. *Journal of Sport and Exercise Psychology* 35: 98-109, 2013.
25. Blazer HJ, Jordan CL, Pederson JA, Rogers RR, Williams TD, Marshall MR, and Ballmann CG. Effects of Time-of-Day Training Preference on Resistance-Exercise Performance. *Research Quarterly for Exercise and Sport*: 1-8, 2020.
26. Borms D, Maenhout A, and Cools AM. Upper quadrant field tests and isokinetic upper limb strength in overhead athletes. *Journal of Athletic Training* 51: 789-796, 2016.
27. Bressel E, Yonker JC, Kras J, and Heath EM. Comparison of static and dynamic balance in female collegiate soccer, basketball, and gymnastics athletes. *Journal of Athletic Training* 42: 42, 2007.
28. Brooks R, Bussiere LF, Jennions MD, and Hunt J. Sinister strategies succeed at the cricket World Cup. *Proceedings of the Royal Society of London Series B: Biological Sciences* 271: S64-S66, 2004.
29. Brophy R, Silvers HJ, Gonzales T, and Mandelbaum BR. Gender influences: the role of leg dominance in ACL injury among soccer players. *British Journal of Sports Medicine* 44: 694-697, 2010.
30. Brown S, Brughelli M, and Cross M. Profiling sprint mechanics by leg preference and position in rugby union athletes. *International Journal of Sports Medicine* 37: 890-897, 2016.
31. Brown SR, Brughelli M, and Lenetsky S. Profiling Single-Leg Balance by Leg Preference and Position in Rugby Union Athletes. *Motor Control* 22: 183-198, 2018.
32. Brown SR, Feldman ER, Cross MR, Helms ER, Marrier B, Samozino P, and Morin J-B. The Potential for a Targeted Strength-Training Program to Decrease Asymmetry and Increase Performance: A Proof of Concept in Sprinting. *International Journal of Sports Physiology and Performance* 12: 1392-1395, 2017.
33. Brumitt J, Nelson K, Duey D, Jeppson M, and Hammer L. Preseason Y Balance Test Scores are not Associated with Noncontact Time-Loss Lower Quadrant Injury in Male Collegiate Basketball Players. *Sports* 7: 4, 2019.
34. Bryson A, Frick B, and Simmons R. The returns to scarce talent: Footedness and player remuneration in European soccer. *Journal of Sports Economics* 14: 606-628, 2013.
35. Buekers M, Ibáñez-Gijón J, Morice AH, Rao G, Mascret N, Laurin J, and Montagne G. Interdisciplinary research: A promising approach to investigate elite performance in sports. *Quest* 69: 65-79, 2017.
36. Bullock GS, Arnold TW, Plisky PJ, and Butler RJ. Basketball players' dynamic performance across competition levels. *The Journal of Strength & Conditioning Research* 32: 3528-3533, 2018.

37. Camomilla V, Bergamini E, Fantozzi S, and Vannozzi G. Trends supporting the in-field use of wearable inertial sensors for sport performance evaluation: A systematic review. *Sensors* 18: 873, 2018.
38. Cardinale M. Commentary on "Towards a Grand Unified Theory of sports performance.". *Human Movement Science* 56: 160-162, 2017.
39. Carey DP, Smith DT, Martin D, Smith G, Skriver J, Rutland A, and Shepherd JW. The bi-pedal ape: Plasticity and asymmetry in footedness. *cortex* 45: 650-661, 2009.
40. Carey DP, Smith G, Smith DT, Shepherd JW, Skriver J, Ord L, and Rutland A. Footedness in world soccer: an analysis of France'98. *Journal of Sports Sciences* 19: 855-864, 2001.
41. Carvalho A, Brown S, and Abade E. Evaluating injury risk in first and second league professional Portuguese soccer: muscular strength and asymmetry. *Journal of Human Kinetics* 51: 19-26, 2016.
42. Castañer M, Andueza J, Hileno R, Puigarnau S, Prat Q, and Camerino O. Profiles of motor laterality in young athletes' performance of complex movements. *Frontiers in Psychology* 9: 916, 2018.
43. Chalmers S, DeBenedictis TA, Zacharia A, Townsley S, Gleeson C, Lynagh M, Townsley A, and Fuller JT. Asymmetry during Functional Movement Screening and injury risk in junior football players: A replication study. *Scandinavian journal of medicine & science in sports* 28: 1281-1287, 2018.
44. Chmielewski TL, Martin C, Lentz TA, Tillman SM, Moser MW, Farmer KW, and Jaric S. Normalization considerations for using the unilateral seated shot put test in rehabilitation. *journal of orthopaedic & sports physical therapy* 44: 518-524, 2014.
45. Clagg SE, Warnock A, and Thomas JS. Kinetic analyses of maximal effort soccer kicks in female collegiate athletes. *Sports Biomechanics* 8: 141-153, 2009.
46. Cohen J. A coefficient of agreement for nominal scales. *Educational and psychological measurement* 20: 37-46, 1960.
47. Corbetta D, Williams J, and Snapp-Childs W. Plasticity in the development of handedness: evidence from normal development and early asymmetric brain injury. *Developmental Psychobiology: The Journal of the International Society for Developmental Psychobiology* 48: 460-471, 2006.
48. Coren S. The lateral preference inventory for measurement of handedness, footedness, eyedness, and earedness: Norms for young adults. *Bulletin of the Psychonomic Society* 31: 1-3, 1993.
49. Cowley JC and Gates DH. Inter-joint coordination changes during and after muscle fatigue. *Human movement science* 56: 109-118, 2017.
50. Croisier J-L, Ganteaume S, Binet J, Genty M, and Ferret J-M. Strength imbalances and prevention of hamstring injury in professional soccer players: a prospective study. *The American journal of sports medicine* 36: 1469-1475, 2008.
51. Cronin JB and Hansen KT. Strength and power predictors of sports speed. *J Strength Cond Res* 19: 349-357, 2005.
52. Čuk T, Leben-Seljak P, and Štefančič M. *Lateral asymmetry of human long bones*. Adam Mickiewicz University, 2001.
53. Daneshjoo A, Rahnama N, Mokhtar AH, and Yusof A. Bilateral and unilateral asymmetries of isokinetic strength and flexibility in male young professional soccer players. *Journal of human kinetics* 36: 45-53, 2013.
54. de Lira CAB, Vargas VZ, Vancini RL, and Andrade MS. Profiling Isokinetic Strength of Shoulder Rotator Muscles in Adolescent Asymptomatic Male Volleyball Players. *Sports* 7: 49, 2019.
55. de Ruijter CJ, de Korte A, Schreven S, and De Haan A. Leg dominance in relation to fast isometric torque production and squat jump height. *European journal of applied physiology* 108: 247, 2010.

56. Donatelli R, Ellenbecker TS, Ekedahl SR, Wilkes JS, Kocher K, and Adam J. Assessment of shoulder strength in professional baseball pitchers. *Journal of Orthopaedic & Sports Physical Therapy* 30: 544-551, 2000.
57. Dos' Santos T, Thomas C, Jones PA, and Comfort P. Asymmetries in isometric force-time characteristics are not detrimental to change of direction speed. *The Journal of Strength & Conditioning Research* 32: 520-527, 2018.
58. Dos'Santos T, Bishop C, Thomas C, Comfort P, and Jones PA. The effect of limb dominance on change of direction biomechanics: A systematic review of its importance for injury risk. *Physical therapy in sport*, 2019.
59. Dos'Santos T, Thomas C, Jones PA, and Comfort P. Asymmetries in single and triple hop are not detrimental to change of direction speed. *Journal of Trainology* 6: 35-41, 2017.
60. Duehring MD, Feldmann CR, and Ebben WP. Strength and conditioning practices of United States high school strength and conditioning coaches. *The Journal of Strength & Conditioning Research* 23: 2188-2203, 2009.
61. Duthie G, Pyne D, and Hooper S. Applied physiology and game analysis of rugby union. *Sports medicine* 33: 973-991, 2003.
62. Ebben WP and Blackard DO. Strength and conditioning practices of National Football League strength and conditioning coaches. *Journal of Strength and Conditioning Research* 15: 48-58, 2001.
63. Ebben WP, Carroll RM, and Simenz CJ. Strength and conditioning practices of National Hockey League strength and conditioning coaches. *The Journal of Strength & Conditioning Research* 18: 889-897, 2004.
64. Ebben WP, Hintz MJ, and Simenz CJ. Strength and conditioning practices of Major League Baseball strength and conditioning coaches. *The Journal of Strength & Conditioning Research* 19: 538-546, 2005.
65. Ellenbecker TS and Mattalino AJ. Concentric isokinetic shoulder internal and external rotation strength in professional baseball pitchers. *Journal of Orthopaedic & Sports Physical Therapy* 25: 323-328, 1997.
66. Ellenbecker TS, Roetert EP, Sueyoshi T, and Riewald S. A descriptive profile of age-specific knee extension flexion strength in elite junior tennis players. *British Journal of Sports Medicine* 41: 728-732, 2007.
67. Ellenbeckert TS. Shoulder internal and external rotation strength and range of motion of highly skilled junior tennis players. *Isokinetics and Exercise Science* 2: 65-72, 1992.
68. Falsone SA, Gross MT, Guskiewicz KM, and Schneider RA. One-arm hop test: reliability and effects of arm dominance. *Journal of Orthopaedic & Sports Physical Therapy* 32: 98-103, 2002.
69. Fleisig GS, Andrews JR, Dillman CJ, and Escamilla RF. Kinetics of baseball pitching with implications about injury mechanisms. *The American journal of sports medicine* 23: 233-239, 1995.
70. Ford KR, Myer GD, and Hewett TE. Valgus knee motion during landing in high school female and male basketball players. *Medicine & Science in Sports & Exercise* 35: 1745-1750, 2003.
71. Fort-Vanmeerhaeghe A, Gual G, Romero-Rodriguez D, and Unnitha V. Lower limb neuromuscular asymmetry in volleyball and basketball players. *Journal of human kinetics* 50: 135-143, 2016.
72. Fort-Vanmeerhaeghe A, Montalvo AM, Sitja-Rabert M, Kiefer AW, and Myer GD. Neuromuscular asymmetries in the lower limbs of elite female youth basketball players and the application of the skillful limb model of comparison. *Physical Therapy in Sport* 16: 317-323, 2015.
73. Fortier A, Turcotte RA, and Pearsall DJ. Skating mechanics of change-of-direction manoeuvres in ice hockey players. *Sports biomechanics* 13: 341-350, 2014.

74. Franceschini KC, Nissola N, Zardo BS, Tadielo GS, and Bonetti LV. Isokinetic performance of shoulder external and internal rotators in adolescent male volleyball athletes. *International Archives of Medicine* 9, 2016.
75. Freitas TT, Martinez-Rodriguez A, Calleja-González J, and Alcaraz PE. Short-term adaptations following Complex Training in team-sports: A meta-analysis. *PLoS One* 12: e0180223, 2017.
76. Gabbard C and Hart S. A question of foot dominance. *The Journal of general psychology* 123: 289-296, 1996.
77. Gabbard SHC. Examining the stabilising characteristics of footedness. *Laterality: Asymmetries of Body, Brain and Cognition* 2: 17-26, 1997.
78. Gkrilias P, Zavvos A, Fousekis K, Billis E, Matzaroglou C, and Tsepis E. Dynamic balance asymmetries in pre-season injury-prevention screening in healthy young soccer players using the Modified Star Excursion Balance Test—a pilot study. *Journal of physical therapy science* 30: 1141-1144, 2018.
79. Gleason BH. Stability of Isometric Strength Asymmetry and Its Relationship to Sprint and Change-of-Direction Performance Asymmetry in Division-I Collegiate Athletes. 2015.
80. Godfrey A, Del Din S, Barry G, Mathers J, and Rochester L. Instrumenting gait with an accelerometer: a system and algorithm examination. *Medical engineering & physics* 37: 400-407, 2015.
81. Gokeler A, Hof A, Arnold M, Dijkstra P, Postema K, and Otten E. Abnormal landing strategies after ACL reconstruction. *Scandinavian journal of medicine & science in sports* 20: e12-e19, 2010.
82. Gonell AC, Romero JAP, and Soler LM. Relationship between the Y balance test scores and soft tissue injury incidence in a soccer team. *International journal of sports physical therapy* 10: 955, 2015.
83. Grealy MA and Mathers JF. Motor control strategies and the effects of fatigue on golf putting performance. *Frontiers in psychology* 4: 1005, 2014.
84. Greenland S and Morgenstern H. Confounding in health research. *Annual review of public health* 22: 189-212, 2001.
85. Greska E, Cortes N, Ringleb S, Onate J, and Van Lunen B. Biomechanical differences related to leg dominance were not found during a cutting task. *Scandinavian journal of medicine & science in sports* 27: 1328-1336, 2017.
86. Grgic J, Lazinica B, Garofolini A, Schoenfeld BJ, Saner NJ, and Mikulic P. The effects of time of day-specific resistance training on adaptations in skeletal muscle hypertrophy and muscle strength: a systematic review and meta-analysis. *Chronobiology international* 36: 449-460, 2019.
87. Gribble PA and Hertel J. Considerations for normalizing measures of the Star Excursion Balance Test. *Measurement in physical education and exercise science* 7: 89-100, 2003.
88. Grouios G, Kollias N, Koidou I, and Poderi A. Excess of mixed-footedness among professional soccer players. *Perceptual and motor skills* 94: 695-699, 2002.
89. Gualdi-Russo E, Rinaldo N, Pasini A, and Zaccagni L. Hand Preference and Performance in Basketball Tasks. *International journal of environmental research and public health* 16: 4336, 2019.
90. Guiard Y. Asymmetric division of labor in human skilled bimanual action: The kinematic chain as a model. *Journal of motor behavior* 19: 486-517, 1987.
91. Hadzic V, Sattler T, Veselko M, Markovic G, and Dervisevic E. Strength asymmetry of the shoulders in elite volleyball players. *Journal of athletic training* 49: 338-344, 2014.
92. Hanna SJ and Abass SA. Comparison in some kinematic variables of layup basketball of older and young players. 2015.
93. Harper DJ, Carling C, and Kiely J. High-Intensity Acceleration and Deceleration Demands in Elite Team Sports Competitive Match Play: A Systematic Review and Meta-Analysis of Observational Studies. *Sports Medicine*: 1-25, 2019.



94. Hart NH, Nimphius S, Cochrane JL, and Newton RU. Leg mass characteristics of Accurate and Inaccurate kickers—an Australian Football perspective. *Journal of Sports Sciences* 31: 1647-1655, 2013.
95. Hart NH, Nimphius S, Spiteri T, and Newton RU. Leg strength and lean mass symmetry influences kicking performance in Australian football. *Journal of sports science & medicine* 13: 157, 2014.
96. Hart NH, Nimphius S, Weber J, Spiteri T, Rantalainen T, Dobbin M, and Newton RU. Musculoskeletal asymmetry in football athletes: a product of limb function over time. *Medicine and science in sports and exercise* 48: 1379-1387, 2016.
97. Hart NH, Spiteri T, Lockie RG, Nimphius S, and Newton RU. Detecting deficits in change of direction performance using the preplanned multidirectional Australian football league agility test. *The Journal of Strength & Conditioning Research* 28: 3552-3556, 2014.
98. Hartley EM, Hoch MC, and Boling MC. Y-balance test performance and BMI are associated with ankle sprain injury in collegiate male athletes. *Journal of science and medicine in sport* 21: 676-680, 2018.
99. Hewett TE, Myer GD, Ford KR, Heidt Jr RS, Colosimo AJ, McLean SG, Van den Bogert AJ, Paterno MV, and Succop P. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *The American journal of sports medicine* 33: 492-501, 2005.
100. Hewitt JK, Cronin JB, and Hume PA. Asymmetry in multi-directional jumping tasks. *Physical Therapy in Sport* 13: 238-242, 2012.
101. Hides J, Fan T, Stanton W, Stanton P, McMahon K, and Wilson S. Psoas and quadratus lumborum muscle asymmetry among elite Australian Football League players. *British Journal of Sports Medicine* 44: 563-567, 2010.
102. Hrysomallis C, McLaughlin P, and Goodman C. Relationship between static and dynamic balance tests among elite Australian Footballers. *Journal of science and medicine in sport* 9: 288-291, 2006.
103. Jäncke L, Peters M, Schlaug G, Posse S, Steinmetz H, and Müller-Gärtner H-W. Differential magnetic resonance signal change in human sensorimotor cortex to finger movements of different rate of the dominant and subdominant hand. *Cognitive Brain Research* 6: 279-284, 1998.
104. Jones PA and Bampouras TM. A comparison of isokinetic and functional methods of assessing bilateral strength imbalance. *The Journal of Strength & Conditioning Research* 24: 1553-1558, 2010.
105. Kobayashi Y, Kubo J, Matsubayashi T, Matsuo A, Kobayashi K, and Ishii N. Relationship between bilateral differences in single-leg jumps and asymmetry in isokinetic knee strength. *Journal of Applied Biomechanics* 29: 61-67, 2013.
106. Kobayashi Y, Kubo J, Matsuo A, Matsubayashi T, Kobayashi K, and Ishii N. Bilateral asymmetry in joint torque during squat exercise performed by long jumpers. *The Journal of Strength & Conditioning Research* 24: 2826-2830, 2010.
107. Korhonen MT, Suominen H, Viitasalo JT, Liikavainio T, Alen M, and Mero AA. Variability and symmetry of force platform variables in maximum-speed running in young and older athletes. *Journal of Applied Biomechanics* 26: 357-366, 2010.
108. Kuki S, Konishi Y, Okudaira M, Yoshida T, Exell T, and Tanigawa S. Asymmetry of force generation and neuromuscular activity during multi-joint isometric exercise. *The Journal of Physical Fitness and Sports Medicine* 8: 37-44, 2019.
109. Kuki S, Konishi Y, Yoshida T, and Tanigawa S. Relationship among Unilateral Stance Isometric Mid-thigh Pull Variables, Sprint Times, and Jump Performance in Collegiate Football Players. *International Journal of Sport and Health Science*: 201832, 2019.

110. Lai WC, Wang D, Chen JB, Vail J, Rugg CM, and Hame SL. Lower quarter Y-balance test scores and lower extremity injury in NCAA division I athletes. *Orthopaedic journal of sports medicine* 5: 2325967117723666, 2017.
111. Landolsi M, Labiadh L, Zarrouk F, Maaref K, Ghannouchi S, Tabka Z, and Lacouture P. Kinematic analysis of the shot-put: A method of assessing the mechanical work of the hand action force. *European journal of sport science* 18: 1208-1216, 2018.
112. Lawler TP and Lawler FH. Left-handedness in professional basketball: prevalence, performance, and survival. *Perceptual and motor skills* 113: 815-824, 2011.
113. Lisman P, Hildebrand E, Nadelen M, and Leppert K. Association of Functional Movement Screen and Y-Balance Test Scores With Injury in High School Athletes. *Journal of strength and conditioning research*, 2019.
114. Lockie RG, Schultz AB, Jeffriess MD, and Callaghan SJ. The relationship between bilateral differences of knee flexor and extensor isokinetic strength and multi-directional speed. *Isokinetics and exercise science* 20: 211-219, 2012.
115. Loffing F, Sölter F, and Hagemann N. Left preference for sport tasks does not necessarily indicate left-handedness: sport-specific lateral preferences, relationship with handedness and implications for laterality research in behavioural sciences. *PLoS one* 9, 2014.
116. Loffing F, Sölter F, Hagemann N, and Strauss B. Accuracy of outcome anticipation, but not gaze behavior, differs against left-and right-handed penalties in team-handball goalkeeping. *Frontiers in psychology* 6: 1820, 2015.
117. Ludwig O, Simon S, Piret J, Becker S, and Marschall F. Differences in the dominant and non-dominant knee valgus angle in junior elite and amateur soccer players after unilateral landing. *Sports* 5: 14, 2017.
118. Luk H-Y, Winter C, O'Neill E, and Thompson BA. Comparison of muscle strength imbalance in powerlifters and jumpers. *The Journal of Strength & Conditioning Research* 28: 23-27, 2014.
119. Madruga-Parera M, Bishop C, Fort-Vanmeerhaeghe A, Beltran-Valls MR, Skok OG, and Romero-Rodríguez D. Interlimb Asymmetries in Youth Tennis Players: Relationships With Performance. *Journal of strength and conditioning research*, 2019.
120. Madruga-Parera M, Bishop C, Read P, and Lake J. Jumping-based Asymmetries are Negatively Associated with Jump, Change of Direction, and Repeated Sprint Performance, but not Linear Speed, in Adolescent Handball Athletes.
121. Malone JJ, Lovell R, Varley MC, and Coutts AJ. Unpacking the black box: applications and considerations for using GPS devices in sport. *International journal of sports physiology and performance* 12: S2-18-S12-26, 2017.
122. Maloney SJ. The relationship between asymmetry and athletic performance: A critical review. *The Journal of Strength & Conditioning Research* 33: 2579-2593, 2019.
123. Maloney SJ, Richards J, Nixon DG, Harvey LJ, and Fletcher IM. Do stiffness and asymmetries predict change of direction performance? *Journal of sports sciences* 35: 547-556, 2017.
124. Maly T, Zahalka F, Bonacin D, Mala L, and Bujnovsky D. Muscular strength and strength asymmetries of high elite female soccer players. *Sport Sci* 8: 7-14, 2015.
125. Maly T, Zahalka F, and Mala L. Unilateral and Ipsilateral Strength Asymmetries in Elite Youth Soccer Players With Respect to Muscle Group and Limb Dominance. *International Journal of Morphology* 34, 2016.
126. Matthew D and Delextrat A. Heart rate, blood lactate concentration, and time–motion analysis of female basketball players during competition. *Journal of sports sciences* 27: 813-821, 2009.
127. McLean J and Ciurczak F. Bimanual dexterity in major league baseball players: a statistical study. *The New England journal of medicine* 307: 1278-1279, 1982.
128. McLean SG, Fellin R, Suedekum N, Calabrese G, Passerallo A, and Joy S. Impact of fatigue on gender-based high-risk landing strategies. *Medicine and science in sports and exercise* 39: 502-514, 2007.

129. Mentiplay BF, Mosler AB, Crossley KM, Carey DL, Sakadjian K, Bodger R, Shipperd B, and Bruder AM. Lower limb musculoskeletal screening in elite female Australian football players. *Physical Therapy in Sport* 40: 33-43, 2019.
130. Mok KM, Bahr R, and Krosshaug T. The effect of overhead target on the lower limb biomechanics during a vertical drop jump test in elite female athletes. *Scandinavian journal of medicine & science in sports* 27: 161-166, 2017.
131. Mulrey C, Ford KR, Shultz SJ, Nguyen A-D, and Taylor JB. Identifying Limb Dominance in Adolescent Female Basketball Players: Implications for Biomechanical Research. Presented at Medicine and science in sports and exercise, 2016.
132. Mulrey CR, Shultz SJ, Ford KR, Nguyen A-D, and Taylor JB. Methods of Identifying Limb Dominance in Adolescent Female Basketball Players: Implications for Clinical and Biomechanical Research. *Clinical journal of sport medicine: official journal of the Canadian Academy of Sport Medicine*, 2018.
133. Mulrey CR, Shultz SJ, Ford KR, Nguyen A-D, and Taylor JB. Methods of Identifying Limb Dominance in Adolescent Female Basketball Players: Implications for Clinical and Biomechanical Research. *Clinical Journal of Sport Medicine* 30: 279-281, 2020.
134. Murray A, Buttfield A, Simpkin A, Sproule J, and Turner AP. Variability of within-step acceleration and daily wellness monitoring in Collegiate American Football. *Journal of science and medicine in sport* 22: 488-493, 2019.
135. Newman MA, Tarpenning KM, and Marino FE. Relationships between isokinetic knee strength, single-sprint performance, and repeated-sprint ability in football players. *Journal of Strength and Conditioning Research* 18: 867-872, 2004.
136. Nightingale SC. A strength and conditioning approach for ice hockey. *Strength & Conditioning Journal* 36: 28-36, 2014.
137. Onate JA, Starkel C, Clifton DR, Best TM, Borchers J, Chaudhari A, Comstock RD, Cortes N, Grooms DR, and Hertel J. Normative functional performance values in high school athletes: The functional pre-participation evaluation project. *Journal of athletic training* 53: 35-42, 2018.
138. Peters M. Footedness: asymmetries in foot preference and skill and neuropsychological assessment of foot movement. *Psychological Bulletin* 103: 179, 1988.
139. Piggott B, Müller S, Chivers P, Papaluca C, and Hoyne G. Is sports science answering the call for interdisciplinary research? A systematic review. *European journal of sport science* 19: 267-286, 2019.
140. Puterman J, Baker J, and Schorer J. Laterality differences in elite ice hockey: An investigation of shooting and catching orientations. *Journal of sports sciences* 28: 1581-1593, 2010.
141. Rahnama N, Lees A, and Bambaecichi E. A comparison of muscle strength and flexibility between the preferred and non-preferred leg in English soccer players. *Ergonomics* 48: 1568-1575, 2005.
142. Read M and Bellamy M. Comparison of hamstring/quadriceps isokinetic strength ratios and power in tennis, squash and track athletes. *British Journal of Sports Medicine* 24: 178-182, 1990.
143. Read PJ, Hughes J, Stewart P, Chavda S, Bishop C, Edwards M, and Turner AN. A needs analysis and field-based testing battery for basketball. *Strength & Conditioning Journal* 36: 13-20, 2014.
144. Read PJ, Oliver JL, De Ste Croix M, Myer GD, and Lloyd RS. A prospective investigation to evaluate risk factors for lower extremity injury risk in male youth soccer players. *Scandinavian journal of medicine & science in sports* 28: 1244-1251, 2018.
145. Riemann BL, Guskiewicz KM, and Shields EW. Relationship between clinical and forceplate measures of postural stability. *Journal of sport rehabilitation* 8: 71-82, 1999.

146. Roell M, Roecker K, Gehring D, Mahler H, and Gollhofer A. Player monitoring in indoor team sports: concurrent validity of inertial measurement units to quantify average and peak acceleration values. *Frontiers in physiology* 9: 141, 2018.
147. Rosene JM, Fogarty TD, and Mahaffey BL. Isokinetic hamstrings: quadriceps ratios in intercollegiate athletes. *Journal of athletic training* 36: 378, 2001.
148. Saccol MF, Gracitelli GC, da Silva RT, de Souza Laurino CF, Fleury AM, dos Santos Andrade M, and da Silva AC. Shoulder functional ratio in elite junior tennis players. *Physical Therapy in Sport* 11: 8-11, 2010.
149. Samadi H, Rajabi R, Minoonejad H, and Aghaiari A. Asymmetries in flexibility, balance and power associated with preferred and non-preferred leg. *World Journal of Sport Sciences* 2: 38-42, 2009.
150. Sannicandro I, Piccinno A, Rosa R, and De Pascalis S. Correlation between functional asymmetry of professional soccer players and sprint. *British Journal of Sports Medicine* 45: 370-371, 2011.
151. Scharoun SM and Bryden PJ. Hand preference, performance abilities, and hand selection in children. *Frontiers in psychology* 5: 82, 2014.
152. Schiltz M, Lehance C, Maquet D, Bury T, Crielaard J-M, and Croisier J-L. Explosive strength imbalances in professional basketball players. *Journal of Athletic Training* 44: 39-47, 2009.
153. Schmitt LC, Paterno MV, Ford KR, Myer GD, and Hewett TE. Strength asymmetry and landing mechanics at return to sport after ACL reconstruction. *Medicine and science in sports and exercise* 47: 1426, 2015.
154. Scott MT, Scott TJ, and Kelly VG. The validity and reliability of global positioning systems in team sport: a brief review. *The Journal of Strength & Conditioning Research* 30: 1470-1490, 2016.
155. Simenz CJ, Dugan CA, and Ebben WP. Strength and conditioning practices of National Basketball Association strength and conditioning coaches. *The Journal of Strength & Conditioning Research* 19: 495-504, 2005.
156. Sinclair J and Hobbs SJ. Bilateral differences in knee and ankle loading of the support limb during maximal instep soccer kicking. *Science & Sports* 31: e73-e78, 2016.
157. Smith CA, Chimera NJ, and Warren M. Association of y balance test reach asymmetry and injury in division I athletes. *Medicine and science in sports and exercise* 47: 136-141, 2015.
158. Spiteri T, Binetti M, Scanlan AT, Dalbo VJ, Dolci F, and Specos C. Physical Determinants of Division 1 Collegiate Basketball, Women's National Basketball League, and Women's National Basketball Association Athletes: With Reference to Lower-Body Sidedness. *The Journal of Strength & Conditioning Research* 33: 159-166, 2019.
159. Steidl-Müller L, Hildebrandt C, Müller E, Fink C, and Raschner C. Limb symmetry index in competitive alpine ski racers: Reference values and injury risk identification according to age-related performance levels. *Journal of sport and health science* 7: 405-415, 2018.
160. Stephens TM, Lawson BR, DeVoe DE, and Reiser RF. Gender and bilateral differences in single-leg countermovement jump performance with comparison to a double-leg jump. *Journal of Applied Biomechanics* 23: 190-202, 2007.
161. Stewart S, Stanton W, Wilson S, and Hides J. Consistency in size and asymmetry of the psoas major muscle among elite footballers. *British journal of sports medicine* 44: 1173-1177, 2010.
162. Stiffler MR, Bell DR, Sanfilippo JL, Hetzel SJ, Pickett KA, and Heiderscheit BC. Star excursion balance test anterior asymmetry is associated with injury status in division I collegiate athletes. *journal of orthopaedic & sports physical therapy* 47: 339-346, 2017.
163. Stöckel T and Carey DP. Laterality effects on performance in team sports: insights from soccer and basketball, in: *Laterality in Sports*. Elsevier, 2016, pp 309-328.

164. Stöckel T and Vater C. Hand preference patterns in expert basketball players: interrelations between basketball-specific and everyday life behavior. *Human movement science* 38: 143-151, 2014.
165. Stöckel T and Weigelt M. Plasticity of human handedness: Decreased one-hand bias and inter-manual performance asymmetry in expert basketball players. *Journal of sports sciences* 30: 1037-1045, 2012.
166. Theoharopoulos A and Tsitskaris G. Isokinetic evaluation of the ankle plantar and dorsiflexion strength to determine the dominant limb in basketball players. *Isokinetics and exercise science* 8: 181-186, 2000.
167. THEOHAROPOULOS A, TSITSKARIS G, and TSAKLIS P. Knee strength of professional basketball players. *The Journal of Strength & Conditioning Research* 14: 457-463, 2000.
168. Thomas C, Comfort P, Dos' Santos T, and Jones PA. Determining bilateral strength imbalances in youth basketball athletes. *International journal of sports medicine* 38: 683-690, 2017.
169. Thomas C, Comfort P, Dos'Santos T, and Jones PA. Determining bilateral strength imbalances in youth basketball athletes. *International journal of sports medicine* 38: 683-690, 2017.
170. Thomas C, Dos'Santos T, Kyriakidou I, Cuthbert M, Fields C, and Jones PA. An investigation into the effect of limb preference on knee mechanics and braking strategy during pivoting in female soccer players: An exploratory study. Presented at The 8th Annual Strength and Conditioning Student Conference; Middlesex University: London, UK, 2017.
171. Thorpe JL and Ebersole KT. Unilateral balance performance in female collegiate soccer athletes. *The Journal of Strength & Conditioning Research* 22: 1429-1433, 2008.
172. Turner E, Munro AG, and Comfort P. Female soccer: Part 1—A needs analysis. *Strength & Conditioning Journal* 35: 51-57, 2013.
173. Tyler TF, Nicholas SJ, Campbell RJ, and McHugh MP. The association of hip strength and flexibility with the incidence of adductor muscle strains in professional ice hockey players. *The American journal of sports medicine* 29: 124-128, 2001.
174. Vaisman A, Guiloff R, Rojas J, Delgado I, Figueroa D, and Calvo R. Lower Limb Symmetry: Comparison of Muscular Power Between Dominant and Nondominant Legs in Healthy Young Adults Associated With Single-Leg-Dominant Sports. *Orthopaedic journal of sports medicine* 5: 2325967117744240, 2017.
175. van Melick N, Meddeler BM, Hoogeboom TJ, Nijhuis-van der Sanden MW, and van Cingel RE. How to determine leg dominance: The agreement between self-reported and observed performance in healthy adults. *PLoS one* 12, 2017.
176. van Santen JA, Pereira C, Sanchez-Santos MT, Cooper C, and Arden NK. Dominant vs. non-dominant hip comparison in bone mineral density in young sporting athletes. *Archives of osteoporosis* 14: 54, 2019.
177. Varley MC, Fairweather IH, and Aughey1, Robert J. Validity and reliability of GPS for measuring instantaneous velocity during acceleration, deceleration, and constant motion. *Journal of sports sciences* 30: 121-127, 2012.
178. Velotta J, Weyer J, Ramirez A, Winstead J, and Bahamonde R. Relationship between leg dominance tests and type of task. Presented at ISBS-Conference Proceedings Archive, 2011.
179. Wang H-K, Macfarlane A, and Cochrane T. Isokinetic performance and shoulder mobility in elite volleyball athletes from the United Kingdom. *British Journal of Sports Medicine* 34: 39-43, 2000.
180. Wang H. Mobility impairment, muscle imbalance, muscle weakness, scapular asymmetry and shoulder injury in elite volleyball athletes. *Journal of sports medicine and physical fitness* 41: 403-410, 2001.
181. Williams JH, Jaskowak DJ, and Tegarden DP. Detecting Structured Variability in the Running Gait Using a GPS-Embedded, Trunk-Mounted Accelerometer. Part 1: Method Description.

182. Wright AA, Dischiavi SL, Smoliga JM, Taylor JB, and Hegedus EJ. Association of Lower Quarter Y-Balance Test with lower extremity injury in NCAA Division 1 athletes: an independent validation study. *Physiotherapy* 103: 231-236, 2017.

**Table 1.** Example physical qualities with proposed tests, metrics and technologies that may be considered for limb dominance assessment in sport.

<b>Physical Quality</b>	<b>Possible Tests</b>	<b>Metrics</b>	<b>Technologies</b>
Strength	Unilateral isometric mid-thigh pull or squat	Peak force, time to peak force, force at different time points	Force plate
Power	Unilateral jumps: drop jump (UDJ), countermovement jump (UCJ), horizontal jump (UHJ)	UDJ: jump height, reactive strength UCMJ: jump height, peak/mean force*, impulse*, reactive strength-modified* UHJ: jump distance, peak/mean force*, resultant force*	Force plate, OptoJump, jump mat or My Jump app
Change of Direction Speed (CODS)	Change-of-direction tasks (CODS): 505, 505-modified	Total time, change of direction deficit	Dual beam electronic timing gates
* indicates that metric can only be computed if a force plate is available.			